

BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email editorial.bmjopen@bmj.com

BMJ Open

Consensus Validation of the POSAMINO (POtentially Serious Alcohol Medication INTERactions in Older adults) Criteria

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-017453
Article Type:	Research
Date Submitted by the Author:	23-Apr-2017
Complete List of Authors:	Holton, Alice; School of Pharmacy Gallagher, Paul; Royal College of Surgeons in Ireland, School of Pharmacy Ryan, Cristin; Royal College of Surgeons in Ireland, Fahey, Tom; Royal College of Surgeons in Ireland, Department of General Practice Cousins, Gráinne; Royal College of Surgeons in Ireland, School of Pharmacy
Primary Subject Heading:	Pharmacology and therapeutics
Secondary Subject Heading:	Health services research, Geriatric medicine
Keywords:	CLINICAL PHARMACOLOGY, Adverse events < THERAPEUTICS, alcohol, GERIATRIC MEDICINE

SCHOLARONE™
Manuscripts

Consensus Validation of the POSAMINO (POtentially Serious Alcohol Medication INteractions in Older adults) Criteria

Ms. Alice E. Holton ¹, Prof. Paul Gallagher¹, Dr. Cristín Ryan¹, Prof. Tom Fahey ² and Dr. Gráinne Cousins¹

Affiliations:

¹School of Pharmacy, Royal College of Surgeons, 1st Floor Ardilaun House Block B, 111 St Stephen’s Green, Dublin 2, Ireland

²HRB Centre for Primary Care Research, Division of Population Health Science, Royal College of Surgeons in Ireland, 123 St. Stephens Green, Dublin 2, Ireland

Corresponding author: Ms. Alice Holton MPSI, RCSI School of Pharmacy, Block B Ardilaun House, Dublin 2. Email: aliceholton@rcsi.ie

Abstract:

Objectives: Older adults are particularly vulnerable to adverse effects from concurrent alcohol and medication use. However, there is limited evidence regarding the prevalence of these adverse outcomes among older adults and there is a lack of consensus regarding what constitutes an alcohol-interactive medicine. The objective of this study was to develop an explicit list of potentially serious alcohol-medication interactions for use in older adults.

Design: Following a systematic review, review of drug compendia and clinical guidance documents, a two-round Delphi consensus method was conducted.

Setting: Ireland and the United Kingdom (UK), primary care and hospital setting.

Participants: The Project Steering Group developed a list of potentially serious alcohol-medication interactions. The Delphi panel consisted of 19 healthcare professionals (general practitioners (GPs), geriatricians, hospital and community pharmacists, clinical pharmacologists and pharmacists and physicians specialising in substance misuse).

Results: An inventory of 52 potentially serious alcohol-medication interactions was developed by the Project Steering Group. British National Formulary (BNF) black dot warnings (n=8) were included in the final criteria as they represent 'potentially serious' interactions. The remaining 44 criteria underwent a two-round Delphi process. In the first round, thirteen criteria were accepted into the POSAMINO criteria. Consensus was not reached on the remaining 31 criteria; nine were removed, and eight additional criteria were included following a review of panellist comments. The remaining 30 criteria went to round-two, with 17 criteria reaching consensus, providing a final list of 38 potentially serious alcohol-medication interactions; central nervous system (n=15), cardiovascular system (n=9), endocrine system (n=5), musculoskeletal system (n=3), infections (n=3), malignant disease & immunosuppression (n=2) and respiratory system (n=1).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Conclusions: POSAMINO is the first set of explicit potentially serious alcohol-medication interactions for use in older adults. These criteria will facilitate the risk stratification of older adults at the point of prescribing.

For peer review only

Strengths and limitations of this study:

- The potentially serious alcohol-medication interactions for use in older adults (POSAMINO) criteria were developed in a robust fashion, using a two-step process involving a systematic review and two-round Delphi process.
- The Delphi consensus technique utilised in this study is flexible and enabled communication from a diverse group of healthcare professionals from both the United Kingdom and Ireland.
- The final criteria will need to be further validated prospectively to quantify the magnitude of risk posed for each criterion for adverse outcomes in older adults.
- Once validated, these criteria have the potential to inform medical decision making and enable healthcare professionals to risk stratify older adults at the point of prescribing and prioritise alcohol screening and brief interventions in high-risk groups.

Background

Population demographics are changing globally, with the proportion and age of the older population continuing to increase. (1) While alcohol consumption tends to decline in older age, recent evidence suggests that drinking occasions tend to become more frequent among older adults. (2) There is also evidence of a cohort effect with successive birth cohorts reporting an increase in alcohol consumption among all age groups, including older adults. (3) Ageing is associated with a variety of physiological changes which may place older adults at an increased risk of alcohol-related health problems. (4, 5) In fact, older adults experience a disproportionate burden of alcohol related-harm; in England between 2009 and 2010, adults aged ≥ 65 years accounted for approximately 44% (461,400) of alcohol-related hospital admissions yet comprised of only 17% of the population.(5) Alcohol-related deaths were also highest among those aged 55 to 74 years. (5)

Polypharmacy is also increasing in older adults. (6) For example, approximately 60% of Irish adults aged ≥ 65 years reported taking five or more medicines in 2012. (6) Certain medicines have the potential to interact with alcohol increasing the risk of medical complications such as hypoglycaemia, hypotension, sedation, gastrointestinal bleeds and liver damage. (7, 8) Alcohol interactive (AI) medicines may interact with alcohol by altering the metabolism (pharmacokinetic) or effects (pharmacodynamic) of alcohol and/or the medication. (8) Interactions may occur with any alcohol or follow a dose response, with the risk or severity of an interaction increasing with increasing levels of alcohol consumption. (7)

While a number of studies have investigated the concurrent use of alcohol and medicines with potential to interact with alcohol among older adults, (9-28) there is a lack of consensus regarding what constitutes an AI medication across studies. Several studies focussed on a wide range of medications, using different drug compendia to identify medications as alcohol interactive, thus leading to a lack of consistency in the inclusion of medicines as alcohol interactive. (9-12, 16, 19, 21-23) Other studies focussed on psychotropic medications alone, specifically sedatives and hypnotics. (13, 15, 17, 18, 20, 24, 26, 28)

While there appears to be a high propensity for alcohol-medication interactions among community dwelling older adults, with between one-in-five to one-in-three older adults potentially susceptible to alcohol-

1 medication interactions, (11, 12, 23) no study to date has examined longitudinal associations of concurrent
2 use with adverse outcomes. An evidence based list of medications which have a significant risk of harm to
3 older patients when combined with alcohol would be useful in a clinical setting, allowing for the
4 identification of older adults whose alcohol consumption places them at increased risk and who would
5 benefit from a preventative intervention. Therefore, the aim of this study is to derive the first set of explicit
6 potentially serious alcohol-medication interactions in older adults.
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Methods

Study design

A Delphi consensus technique was used to develop the list of potentially serious alcohol-medication interactions in older adults. The Delphi method allows a consensus opinion to be reached amongst a panel of experts through an iterative process of questionnaires. (29) Ethics approval for this study was obtained from the Royal College of Surgeons in Ireland (Reference Number REC1097).

Compilation of initial list of potentially serious alcohol-medication interactions

An extensive list of medications with potential to interact with alcohol was identified by the Project Steering Group following a comprehensive systematic search using MEDLINE (PubMed), Embase, Scopus and Web of Science databases. A combination of the following keywords and MeSH terms were used: “ethanol”, “alcohol”, “drug interactions”, “drug alcohol interaction” and “aged”. This search was supplemented by a search in Google Scholar and by hand searching references of retrieved articles. The search was restricted to English language articles and articles published since January 1990. Furthermore, the British National Formulary (BNF), Stockley’s Drug Interactions and Martindale Complete Drug Reference drug compendia were also searched (30-32). Additional documents such as clinical guidance documents (33) and previous reviews (7, 8) were also accessed. Information extracted included: medication name/class, potential adverse outcome(s), whether an interaction is likely to occur with any alcohol consumption or with heavy consumption using national low risk drinking guidelines (34, 35) and, if reported, evidence supporting the interaction. The list of medications was organised according to the BNF physiological classification system. There was considerable heterogeneity across reference sources in terms of identifying medications as having potential to interact with alcohol, with no age specific information for interactions. Furthermore, there were inconsistencies in relation to the quantity of concurrent alcohol consumption that should be avoided. For example, the Martindale mentioned that alcohol combined with NSAIDs may increase the risk of gastrointestinal bleeds with no mention of alcohol consumption patterns, Stockley’s described that both NSAIDs and excessive alcohol use carry the risk of gastrointestinal adverse effects, while NSAIDs and alcohol were not included as an interaction in the BNF. For the next step, the Project Steering Group

reviewed and assessed each medication or drug class. Using the definition of “necessary to avoid” previously described by Dreischulte et al. (36), it was considered necessary to avoid concurrent alcohol with a specific medication if, in an average older adult concurrent use of alcohol with this medication would be considered unsafe because (a) there is sufficient evidence that the patient is likely to be harmed, and (b) the likely harm to the patient is large enough to be clinically significant.

Drugs were excluded for the following reasons: their interaction with alcohol was not considered serious; the medication is not licensed for use in older adults; the medication is only administered in a hospital environment (for example Trabectedin), interaction with alcohol may only occur during alcohol withdrawal or has been withdrawn from the market in the UK or Ireland. Following these exclusions a truncated list was further reviewed by the Project Steering Group by a consensus discussion. In this process the Project Steering Group agreed, that BNF black dot warnings would be included in the final list as these are potentially serious interactions and concomitant administration of alcohol should be avoided. (30) Furthermore, the amount of alcohol to be avoided with each medication class was defined through group discussion, based on both the evidence available and the members’ own clinical experience. Additionally, it was agreed that the following adverse outcomes associated with concurrent use were defined using clinical guidance documents and reviews to ensure clarity among all participants: orthostatic hypotension (37), hypoglycaemia (38) and lactic acidosis (39).

Selection of the Delphi panel

A total of 51 experts from the Republic of Ireland and United Kingdom, were invited (via e-mail or letter) to participate as part of the Delphi consensus panel. Participants were identified by the Project Steering Group on the basis of their clinical expertise or knowledge of alcohol-medication interactions or care of older persons. Reasons for non-participation were not required; however in some instances they were provided, they included time commitments and lack of clinical knowledge of alcohol-medication interactions. In total, 19 participants (37%) agreed to participate and written consent was received from all participants before the study. The panel consisted of general practitioners (GPs) (n=5), geriatricians (n=3), hospital pharmacists with expertise in care of older adults (n=3), community pharmacists (n=3), clinical pharmacologists (n=2),

clinical pharmacists specialising in substance misuse (n=2) and a physician specialising in substance misuse (n=1).

The Delphi validation technique and process

An online questionnaire was piloted by two pharmacists and two GPs, to identify any potential problems and to approximate the completion time for the survey. Following amendments, the 19 participants were sent a link to the online POSAMINO questionnaire (via SurveyGizmo®) in March 2016. Participants were given four weeks to complete the online questionnaire and all participants were sent a reminder email after 2 weeks to encourage participation. The questionnaire was both anonymous and confidential.

The quantity of alcohol per standard drink (SD) (10 grammes alcohol in Ireland) or unit (8 grammes alcohol in the United Kingdom), was defined at the beginning of the survey. As some interactions are listed with any alcohol consumption and others with ‘heavy’ consumption, we provided definitions of both at the beginning of the questionnaire. Any alcohol consumption was defined as ≥ 1 SD of alcohol in Ireland or ≥ 1 unit of alcohol in the UK, with heavy consumption defined according to Irish and UK National low risk drinking guidelines. (34, 35) The panel were instructed to evaluate each potential interaction listed according to the “necessary to avoid” framework described above. (36) Medications were categorised according to the BNF physiological classification system. Each statement was presented in the same format stating: It is necessary to avoid [quantity of alcohol] (any or heavy) with [medications/ drug class], followed by a brief rationale for the statement. For example: *It is necessary to avoid: Any alcohol consumption with first generation anti-histamines (for example promethazine). (Rationale: Concurrent alcohol consumption with first generation anti-histamines can increase the risk of sedation).*

Participants were asked to rate their agreement with each statement, using a 5-point Likert scale (1-strongly disagree to 5-strongly agree) along with any additional comments, or suggestions for additional medicines to be included.

After each round, the median and interquartile ranges (IQR) were calculated for each statement. Consistent with previous Delphi consensus studies, the consensus level required for a statement to be retained was defined *a priori* as a median of 4 or 5 with a lower quartile value of ≥ 4 . (40) If a statement had an upper

quartile value of ≤ 2 , this indicated there was general disagreement with the statement between panel members, and the statement was rejected. If group consensus was not reached, the criteria were reviewed by the Project Steering Group and were removed based on comments or revised and included in the second questionnaire. In the second questionnaire, panellists were provided with links to the most recent evidence relating to each of the alcohol-medication interactions, to help facilitate their decisions due to uncertainty in round 1. As with round one, the median response and IQR were calculated and evaluated by the Project Steering Group using the same thresholds to determine consensus between the panel members. If consensus was not reached following the second round, the statement was rejected. Statistical analysis was performed using STATA version 13 and Microsoft Excel 2010.

Results

Following an initial review of the literature, Anatomical Therapeutic Chemical (ATC) codes for a total of 364 AI medicines were extracted by the Project Steering Group. Medicines were classified according to drug classes and organised according to BNF physiological classification systems. From this list, a total of 63 statements were initially compiled based on the ‘necessary to avoid’ framework. (36) Following group discussions, 11 statements were removed/updated or merged together, based on the steering groups clinical knowledge or experience with medicines. For example, the steering group decided to no longer classify benzodiazepines according to the duration of action, as both long acting and short acting have the potential to interact with alcohol. Furthermore, a total of eight BNF black dot warnings were included in the final criteria as they represent potentially serious interactions.

All 19 panel members completed the round one questionnaire, in which forty-four statements were presented. Consensus was reached on 13 statements, with no statements rejected (**Table 1**). Thirty one statements were reviewed by the steering group, with nine statements removed based on comments from the panel, if the interaction was not relevant to older adults or the interaction was not of clinical significance for example: heavy alcohol consumption with vitamin A preparations. Furthermore, eight new statements were included in round two based on comments from panellists at the first round.

A total of 18 out of the 19 recruited participants completed the round two questionnaire. Of the 30 statements included in the second questionnaire, consensus was reached for 17 statements. The remaining 13 statements were rejected as no consensus was reached. In total, consensus was reached for 30 potentially serious drug alcohol interactions in older adults, with the inclusion of BNF black dot warnings the final list was 38 statements.

INSERT TABLE 1 HERE

This final 38 item POSAMINO criteria (**Table 2**) were organised over the following physiological systems:

Central nervous system (n=15), cardiovascular system (n=9), endocrine system (n=5), musculoskeletal system (n=3), infections (n=3), malignant disease and immunosuppression (n=2) and respiratory system (n=1).

INSERT TABLE 2 HERE

Discussion

While older adults experience a disproportionate burden of alcohol related harm, (5) research suggests that healthcare professionals are less likely to discuss alcohol consumption with older adults. (41, 42) Flagging older adults at the point of prescribing an alcohol-interactive medication may facilitate targeted screening and interventions to help reduce harm. Despite the high propensity for alcohol-medication interactions among community dwelling older adults, (9-28) there is still a lack of consensus regarding what constitutes an AI medication.

This study reports the development of a set of 38 explicit criteria, POSAMINO, for identifying older adults at risk of potentially serious alcohol-medication interactions. The development of these criteria is important for several reasons. In the absence of an explicit list of alcohol-interactive medications, multiple drug reference sources have been used in previous studies leading to a lack of consistency in the inclusion of medications across studies, (11, 12, 19, 22, 23) which may result in biased estimates of potential risk among older adults. Furthermore, some of these interactions may be theoretical with trivial clinical significance. Therefore, this study developed a consensus-based set of explicit criteria for potentially serious alcohol-medication interactions in older adults rather than an exhaustive list of medications with potential to interact with alcohol.

Principle findings

The final POSAMINO criteria consist of seven different drug classes. Central nervous system (CNS) medicines represent 40% of the criteria. Estimates from our previous study indicate that approximately one in five older Irish adults are exposed to CNS agents, with over half of patients using CNS agents reporting concurrent alcohol consumption. (12) Nine of the final 38 criteria concern cardiovascular agents, another widely used drug class among aged adults. (12) There is also a high-risk of adverse effects associated with these agents, with anti-platelets, diuretics and anticoagulants identified as the most common drug classes involved in preventable drug-related admissions. (43) Analgesics such as non-steroidal anti-inflammatory drugs (NSAIDs), opioids, paracetamol and gabapentin (used for neuropathic pain) were also included in the final POSAMINO criteria. (43) The prevalence of chronic pain increases with age and older adults may

consume alcohol to help cope and manage their pain. (45) This behaviour may also predispose older adults to adverse outcomes associated with concurrent use with analgesics (for example increased sedation, falls and gastrointestinal bleeds).

Clinical implications and future research

This study adds to a growing body of research investigating the concurrent use of alcohol and alcohol interactive medicines in older adults. Undoubtedly, adverse drug events (ADEs) represent a major burden on healthcare, with ADEs detected in 26.3% of patients aged ≥ 65 years admitted to an Irish hospital with acute illnesses. (44) The increasing use of multiple medications, (6) combined with an increased frequency of alcohol consumption (2) and age-related physiological changes, may predispose older adults to these adverse events. The significant burden of alcohol-related harm and mortality among older adults, (5) indicates a pressing need for future interventions to minimise risk. The final criteria developed from this study, will need to be further validated prospectively to quantify the magnitude of risk posed for each criterion for adverse outcomes in older adults. Once validated, these criteria have the potential to inform medical decision making and enable healthcare professionals to risk stratify older adults at the point of prescribing and prioritise alcohol screening and brief interventions in high-risk groups. This is essential, as older people have previously reported little knowledge about risks associated with the concurrent use of alcohol and medications. (45)

Strengths and Limitations

The criteria were developed in a robust fashion, using a two-step process involving a systematic review and two-round Delphi process. The Delphi technique is flexible and for this study enabled communication from a group of nineteen healthcare professionals from both the United Kingdom and Ireland. Eighteen of the 19 participants completed the two rounds, with participants providing numerous comments and suggestions in both rounds. Similar to previous studies, participants remained anonymous and were not provided with feedback following the first round in order to remove the risk of potential bias. (40)

Inevitably, there were limitations to this Delphi study. With all Delphi studies, participants' judgements may be deemed subjective. In order to reduce this potential bias, a diverse group of healthcare professionals with

expertise or interest in the care of older adults were selected. With explicit criteria, there is also a need for regular updates and revision due to the availability of more up to date information after development. (46) The developed POSAMINO criteria do not apply to older adults diagnosed with an alcohol use disorder, as chronic heavy alcohol consumption can substantially increase the activity of the cytochrome P450 metabolising enzyme CYP2E1. (8) Finally, older adults may also experience chronic illnesses which affect the metabolism of both alcohol and medications. As a result, it is important that healthcare professionals also consider these comorbid diseases when assessing the risk for potential adverse outcomes. (47)

Conclusions

Using a systematic review and a two-round Delphi consensus method, we have developed the first set of explicit potentially serious alcohol-medication interactions (POSAMINO) for use in older adults. These criteria will allow for the risk stratification of older adults at the point of prescribing, and prioritise alcohol screening and brief alcohol interventions in high-risk groups.

Declarations:

Authors' contributions: AH, PG, CR, TF & GC (the project steering group) conceived and designed this study. AH & GC conducted initial literature search. AH communicated with Delphi participants. AH analysed the data after each round. All Project Steering Group members were involved in interpretation of the data. AH & GC drafted the manuscript. All co-authors revised the manuscript and have given the approval for publication.

Competing interests: The authors declare that they have no competing interests.

Ethics approval: Ethical approval for this study was approved by the Royal College of Surgeons in Ireland (RCSI) ethics committee (REC 1097).

Acknowledgements: We would like to thank all Delphi participants who contributed to the development of the POSAMINO criteria.

Funding: This study was funded by the Royal College of Surgeons in Ireland (RCSI), School of Pharmacy, Dublin 2, Ireland.

Data sharing statement: Additional data is available by request from the corresponding author.

References

1. United Nations, Department of Economic and Social Affairs PD. World Population Prospects: The 2015 Revision, Key Findings and Advance Tables. New York: 2015. Available from: https://esa.un.org/unpd/wpp/Publications/Files/Key_Findings_WPP_2015.pdf

2. Britton A, Ben-Shlomo Y, Benzeval M, Kuh D, Bell S. Life course trajectories of alcohol consumption in the United Kingdom using longitudinal data from nine cohort studies. *BMC Med.* 2015;13:47.

3. Meng Y, Holmes J, Hill-McManus D, Brennan A, Meier PS. Trend analysis and modelling of gender-specific age, period and birth cohort effects on alcohol abstinence and consumption level for drinkers in Great Britain using the General Lifestyle Survey 1984-2009. *Addiction.* 2014;109(2):206-15.

4. Huang AR, Mallet L, Rochefort CM, Egual T, Buckeridge DL, Tamblyn R. Medication-related falls in the elderly: causative factors and preventive strategies. *Drugs Aging.* 2012;29(5):359-76.

5. Wadd S, Papadopoulos C. Drinking behaviour and alcohol-related harm amongst older adults: analysis of existing UK datasets. *BMC Res. Notes.* 2014;7:741.

6. Moriarty F, Hardy C, Bennett K, Smith SM, Fahey T. Trends and interaction of polypharmacy and potentially inappropriate prescribing in primary care over 15 years in Ireland: a repeated cross-sectional study. *BMJ open.* 2015;5(9):e008656.

7. Moore AA, Whiteman EJ, Ward KT. Risks of combined alcohol/medication use in older adults. *Am. J. Geriatr. Pharmacother.* 2007;5(1):64-74.

8. Weathermon R, Crabb DW. Alcohol and medication interactions. *Alcohol Res Health.* 1999;23(1):40-54.

9. Adams WL. Potential for adverse drug-alcohol interactions among retirement community residents. *J. Am. Geriatr. Soc.* 1995;43(9):1021-5.

10. Aira M, Hartikainen S, Sulkava R. Community prevalence of alcohol use and concomitant use of medication - A source of possible risk in the elderly aged 75 and older? *Int. J. Geriatr. Psychiatry.* 2005;20(7):680-5.

11. Breslow RA, Dong C, White A. Prevalence of alcohol-interactive prescription medication use among current drinkers: United States, 1999 to 2010. *Alcohol. Clin. Exp. Res.* 2015;39(2):371-9.

12. Cousins G, Galvin R, Flood M, Kennedy M-C, Motterlini N, Henman MC, et al. Potential for alcohol and drug interactions in older adults: evidence from the Irish longitudinal study on ageing. *BMC Geriatr.* 2014;14.

13. Del Rio MC, Prada C, Alvarez FJ. Do Spanish patients drink alcohol while undergoing treatment with benzodiazepines? *Alcohol.* 2002;26(1):31-4.

14. Del Río MC, Prada C, Alvarez FJ. The use of medication and alcohol among the Spanish population. *Br. J. Clin. Pharmacol.* 1996;41(3):253-5.

15. Du Y, Scheidt-Nave C, Knopf H. Use of psychotropic drugs and alcohol among non-institutionalised elderly adults in Germany. *Pharmacopsychiatry.* 2008;41(6):242-51.

16. Forster LE, Pollow R, Stoller EP. Alcohol use and potential risk for alcohol-related adverse drug reactions among community-based elderly. *J. Community Health*. 1993;18(4):225-39.
17. Ilomaki J, Korhonen MJ, Enlund H, Hartzema AG, Kauhanen J. Risk drinking behavior among psychotropic drug users in an aging Finnish population: the FinDrink study. *Alcohol*. 2008;42(4):261-7.
18. Ilomaeki J, Gnjdjic D, Hilmer SN, Le Couteur DG, Naganathan V, Cumming RG, et al. Psychotropic drug use and alcohol drinking in community-dwelling older Australian men: the CHAMP study. *Drug and Alcohol Review*. 2013;32(2):218-22.
19. Immonen S, Pitkälä K. The prevalence of potential alcohol-drug interactions in older adults. *Eur. Geriatr. Med*. 2012;3:S134.
20. John U, Baumeister SE, Volzke H, Meyer C, Ulbricht S, Alte D. Sedative, hypnotic, anxiolytic and opioid medicament use and its co-occurrence with tobacco smoking and alcohol risk drinking in a community sample. *BMC Public Health*. 2007;7:337.
21. Onder G, Landi F, Della Vedova C, Atkinson H, Pedone C, Cesari M, et al. Moderate alcohol consumption and adverse drug reactions among older adults. *Pharmacoepidemiol. Drug Saf*. 2002;11(5):385-92.
22. Pringle KE, Ahern FM, Heller DA, Gold CH, Brown TV. Potential for alcohol and prescription drug interactions in older people. *J. Am. Geriatr. Soc*. 2005;53(11):1930-6.
23. Qato DM, Manzoor BS, Lee TA. Drug-Alcohol Interactions in Older US Adults. *J. Am. Geriatr. Soc*. 2015;63(11):2324-31.
24. Sheahan SL, Coons SJ, Robbins CA, Martin SS, Hendricks J, Latimer M. Psychoactive medication, alcohol use, and falls among older adults. *J. Behav. Med*. 1995;18(2):127-40.
25. Swift W, Stollznow N, Pirota M. The use of alcohol and medicines among Australian adults. *Aust. N. Z. J. Public Health*. 2007;31(6):529-32.
26. Veldhuizen S, Wade TJ, Cairney J. Alcohol consumption among Canadians taking benzodiazepines and related drugs. *Pharmacoepidemiol. Drug Saf*. 2009;18(3):203-10.
27. Wong H, Heuberger R, Logomarsino J, Hewlings S. Associations between alcohol use, polypharmacy and falls in older adults. *Nurs. Older People*. 2016;28(1):30-6.
28. Lagnaoui R, Moore N, Dartigues JF, Fourrier A, Begaud B. Benzodiazepine use and wine consumption in the French elderly. *Br. J. Clin. Pharmacol*. 2001;52(4):455-6.
29. Hsu C-C, Sandford BA. The Delphi technique: making sense of consensus. *Practical assessment, research & evaluation*. 12(10):1-8.
30. Joint Formulary Committee (2015) British National Formulary (BNF). 69th Ed., London- British Medical Association and Royal Pharmaceutical Society of Great Britain.
31. Sweetman S. Martindale: The Complete Drug Reference London: Pharmaceutical Press; [cited 2015]. Available from: <https://www.medicinescomplete-com.proxy.library.rcsi.ie/mc/martindale/current/index.htm>.
32. Stockley's Drug Interactions [online] [Internet]. Pharmaceutical Press. Available from: <https://www.medicinescomplete.com/mc/stockley/current/>.

33. National Institute on Alcohol Abuse and Alcoholism (NIAAA). Harmful Interactions mixing alcohol with medicines. 2014. Available from: https://pubs.niaaa.nih.gov/publications/medicine/Harmful_Interactions.pdf

34. Hope. A Standard Drink in Ireland: What strength? 2009 Health Service Executive- Alcohol Implementation Group

35. UK Chief Medical Officers' Alcohol Guidelines Review: Summary of the proposed new guidelines. Department of Health January 2016 Available from: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/489795/summary.pdf

36. Dreischulte T, Grant AM, McCowan C, McAnaw JJ, Guthrie B. Quality and safety of medication use in primary care: consensus validation of a new set of explicit medication assessment criteria and prioritisation of topics for improvement. *BMC Clin. Pharmacol.* 2012;12:5.

37. LANIER JB, MOTE MB, CLAY EC. Evaluation and Management of Orthostatic Hypotension. *Am. Fam. Physician.* 2011;84(5):527-36.

38. European Medicines Agency (EMA). Guideline on clinical investigation of medicinal products in the treatment or prevention of diabetes mellitus Committee for Medicinal Products for Human Use , 2012 Available from: http://www.ema.europa.eu/docs/en_GB/document_library/Scientific_guideline/2012/06/WC500129256.pdf

39. Salpeter SR, Greyber E, Pasternak GA, Salpeter EE. Risk of fatal and nonfatal lactic acidosis with metformin use in type 2 diabetes mellitus: Systematic review and meta-analysis. *Arch. Intern. Med.* 2003;163(21):2594-602.

40. Cooper JA, Ryan C, Smith SM, Wallace E, Bennett K, Cahir C, et al. The development of the PROMPT (PRescribing Optimally in Middle-aged People's Treatments) criteria. *BMC Health Serv Res.* 2014;14:484.

41. Duru OK, Xu H, Tseng C-H, Mirkin M, Ang A, Tallen L, et al. Correlates of Alcohol-Related Discussions between Older Adults and Their Physicians. *J. Am. Geriatr. Soc.* 2010;58(12):2369-74.

42. Kuerbis A, Sacco P, Blazer DG, Moore AA. Substance Abuse Among Older Adults. *Clin. Geriatr. Med.* 2014;30(3):629-54.

43. Howard RL, Avery AJ, Slavenburg S, Royal S, Pipe G, Lucassen P, et al. Which drugs cause preventable admissions to hospital? A systematic review. *Br. J. Clin. Pharmacol.* 2007;63(2):136-47.

44. Hamilton H, Gallagher P, Ryan C, Byrne S, O'Mahony D. Potentially inappropriate medications defined by stopp criteria and the risk of adverse drug events in older hospitalized patients. *Arch. Intern. Med.* 2011;171(11):1013-9.

45. Zanjani F, Hoogland AI, Downer BG. Alcohol and prescription drug safety in older adults. *Drug, healthcare and patient safety.* 2013;5:13-27.

46. O'Mahony D, O'Sullivan D, Byrne S, O'Connor MN, Ryan C, Gallagher P. STOPP/START criteria for potentially inappropriate prescribing in older people: version 2. *Age Ageing.* 2015;44(2):213-8.

47. Slattum PW, Hassan OE. Medications, Alcohol, and Aging. In: Kuerbis A, Moore AA, Sacco P, Zanjani F, editors. *Alcohol and Aging: Clinical and Public Health Perspectives.* Cham: Springer International Publishing; 2016. p. 117-129.

Table 1: Results from the 2-round Delphi process

	Potentially serious interactions as listed in the BNF	Round 1					Round 2					Final Criteria
		Total	Accept	Revise	Reject	Removal	New statements included	Total	Accept	Revise	Reject	
Physiological System												
Cardiovascular System	1	14	1	13	0	3	2	12	7	0	5	9
Respiratory System	0	1	1	0	0	0	0	0	0	0	0	1
Central Nervous System	4	12	6	6	0	1	5	10	5	0	5	15
Infections	2	5	0	5	0	3	0	2	1	0	1	3
Endocrine System	0	5	3	2	0	0	0	2	2	0	0	5
Obstetrics, Gynaecology, and Urinary Tract Disorders	0	2	0	2	0	1	1	2	0	0	2	0
Malignant Disease and Immunosuppression	1	1	0	1	0	0	0	1	1	0	0	2
Nutrition and Blood	0	1	0	1	0	1	0	0	0	0	0	0
Musculoskeletal System	0	3	2	1	0	0	0	1	1	0	0	3
Total	8	44	13	31	0	9	8	30	17	0	13	38

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 2: Final POSAMINO Criteria (n=38)

CVS System	
1.	Heavy alcohol consumption with multiple anti-hypertensive combinations <i>Rationale: Concurrent use of alcohol consumption and anti-hypertensives may increase the risk of orthostatic hypotension</i>
2.	Heavy alcohol consumption with warfarin (& phenindione) <i>Rationale: Heavy episodic alcohol consumption is associated with an increased risk of major bleeds</i>
3.	Heavy alcohol consumption with regular use of low dose aspirin (75mg) <i>Rationale: Heavy alcohol consumption combined with aspirin may cause a small increase in gastrointestinal blood loss</i>
4.	Heavy alcohol consumption with both regular and as required nitrates (e.g. glyceryl trinitrate, isosorbide dinitrate and isosorbide mononitrate) <i>Rationale: The combined haemodynamic effects of alcohol and nitrates, increases the risk of exaggerated hypotension.</i>
5.	Heavy alcohol consumption with the vasodilatory medication nicorandil <i>Rationale: The combined haemodynamic effects of alcohol & nicorandil; increases the risk of exaggerated hypotension.</i>
6.	Heavy alcohol consumption with the combined use of both nitrates and vasodilator medication (e.g. nicorandil) <i>Rationale: The combined haemodynamic effects of alcohol and nitrates, increases the risk of exaggerated hypotension.</i>
7.	Heavy alcohol consumption with diuretics (e.g. loop diuretics (furosemide), thiazide diuretics (bendroflumethiazide) & potassium sparing diuretics (amiloride)) <i>Rationale: The concurrent use of alcohol and anti-hypertensives may increase the risk of orthostatic hypotension.</i>
8.	Heavy alcohol consumption with alpha blockers (e.g. terazosin) <i>Rationale: The concurrent use of alcohol and anti-hypertensives may increase the risk of orthostatic hypotension.</i>
9.	Heavy alcohol consumption with centrally acting anti-hypertensives (e.g. clonidine or methyldopa) <i>Rationale: Alcohol consumption combined with centrally acting anti-hypertensives may increase the risk of sedation.</i>
Respiratory System	
1.	Any alcohol consumption with first generation anti-histamines (e.g. promethazine) <i>Rationale: Concurrent alcohol consumption with first generation anti-histamines can increase the risk of sedation</i>
Central Nervous System	
1.	Heavy alcohol consumption with benzodiazepines (e.g. diazepam) and benzodiazepine related medications (e.g. zopiclone) <i>Rationale: Alcohol consumption combined with benzodiazepines and benzodiazepine related medications may enhance CNS depressant effects</i>
2.	Heavy alcohol consumption combined with opioids <i>Rationale: Alcohol consumption combined with opioids may enhance CNS depressant effects of alcohol</i>
3.	Heavy alcohol consumption with duloxetine <i>Rationale: Heavy alcohol consumption combined with duloxetine may increase the risk of hepatotoxicity</i>
4.	Heavy alcohol consumption with all anti-psychotics <i>Rationale: Alcohol consumption combined with antipsychotics may increase the risk of sedation</i>
5.	Any alcohol consumption with barbiturates <i>Rationale: Alcohol consumption combined with barbiturates may increase the risk of sedation.</i>
6.	Heavy alcohol consumption with anti-epileptic drugs (AEDs) <i>Rationale: Heavy alcohol consumption can increase the risk of seizures and sedation in patients taking AEDs.</i>
7.	Any alcohol consumption with tricyclic anti-depressants (TCAs) <i>Rationale: Alcohol consumption combined with antidepressants may enhance the CNS depressant effects of alcohol</i>
8.	Any alcohol consumption with tetracyclic antidepressants <i>Rationale: Alcohol consumption combined with antidepressants may enhance the CNS depressant effects of alcohol</i>
9.	Any alcohol consumption with mirtazapine <i>Rationale: Alcohol consumption combined with antidepressants may enhance the CNS depressant effects of alcohol</i>
10.	Any alcohol consumption with monoamine oxidase inhibitors (MAOIs) <i>Rationale: A potentially life-threatening hypertensive reaction can develop in patients taking non-selective MAOIs who consume drinks rich in tyramine (e.g. wines, beers and non-alcoholic beers)</i>
11.	Heavy alcohol consumption with long term regular paracetamol use (e.g. 1g four times a day) <i>Rationale: Heavy alcohol consumption may increase the risk of paracetamol hepatotoxicity especially if alcohol intake is abruptly stopped.</i>
12.	Heavy alcohol consumption with gabapentin (when used for neuropathic pain) <i>Rationale: Alcohol combined with medications for neuropathic pain may increase the risk of sedation</i>
13.	Heavy alcohol consumption with pramipexole or amantadine <i>Rationale: Alcohol may increase the risk of sedation</i>
14.	Heavy alcohol consumption with apomorphine

Rationale: Alcohol combined with apomorphine may increase the risk of orthostatic hypotension

15. Heavy alcohol consumption with levodopa (alone or in combination with carbidopa)

Rationale: Alcohol combined with levodopa (alone or in combination with carbidopa) may increase the risk of orthostatic hypotension.

Endocrine

1. Heavy alcohol consumption with insulin

Rationale: Alcohol consumption may enhance the hypoglycaemic effect of insulin.

2. Heavy alcohol consumption with metformin

Rationale: Heavy alcohol consumption combined with metformin may increase the risk of lactic acidosis

3. Heavy alcohol consumption with sulphonylureas

Rationale: Alcohol consumption can enhance the hypoglycaemic effects of antidiabetics

4. Heavy alcohol consumption with meglitinides (e.g. nateglinide)

Rationale: Alcohol consumption can enhance the hypoglycaemic effects of antidiabetics

5. Heavy alcohol consumption with thiazolidinediones (e.g. pioglitazone)

Rationale: Alcohol consumption can enhance the hypoglycaemic effects of antidiabetics

Musculoskeletal and joint diseases

1. Heavy alcohol consumption with any nonsteroidal anti-inflammatory drugs NSAIDs (including COX-2 inhibitors)

Rationale: Heavy alcohol consumption and NSAID use carry an increased risk of gastrointestinal bleeds

2. Heavy alcohol consumption combined with methotrexate or leflunomide

Rationale: Heavy alcohol consumption combined with methotrexate or leflunomide may increase the risk of hepatotoxicity

3. Heavy alcohol consumption with oral muscle relaxants (e.g. baclofen)

Rationale: Concurrent alcohol consumption and muscle relaxants can increase the risk of CNS depression.

Malignant disease and immunosuppression

1. Any alcohol consumption with procarbazine

Rationale: A disulfiram-like reaction can occur when alcohol is given with procarbazine

2. Heavy alcohol consumption with interferon alpha or interferon beta

Rationale: Heavy alcohol consumption combined with interferons may increase the risk of hepatotoxicity and reduce the response to treatment with interferon.

Infections

1. Heavy alcohol consumption with anti-mycobacterial medications such as isoniazid, pyrazinamide, ethionamide and rifampicin (alone or in combination)

Rationale: Alcohol combined with anti-mycobacterial medications increase the risk of hepatotoxicity

2. Any alcohol consumption with cycloserine

Rationale: Alcohol consumption may increase the risk of seizures in patients taking cycloserine.

3. Any alcohol consumption with metronidazole or tinidazole

Rationale: A disulfiram-like reaction can occur when alcohol is given with metronidazole

BMJ Open

Consensus Validation of the POSAMINO (POtentially Serious Alcohol Medication INTERactions in Older adults) Criteria

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-017453.R1
Article Type:	Research
Date Submitted by the Author:	01-Aug-2017
Complete List of Authors:	Holton, Alice; School of Pharmacy Gallagher, Paul; Royal College of Surgeons in Ireland, School of Pharmacy Ryan, Cristin; Royal College of Surgeons in Ireland, Fahey, Tom; Royal College of Surgeons in Ireland, Department of General Practice Cousins, Gráinne; Royal College of Surgeons in Ireland, School of Pharmacy
Primary Subject Heading:	Pharmacology and therapeutics
Secondary Subject Heading:	Health services research, Geriatric medicine
Keywords:	CLINICAL PHARMACOLOGY, Adverse events < THERAPEUTICS, alcohol, GERIATRIC MEDICINE

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Consensus Validation of the POSAMINO (POtentially Serious Alcohol Medication INteractions in Older adults) Criteria

Ms. Alice E. Holton ¹, Prof. Paul Gallagher¹, Dr. Cristín Ryan¹, Prof. Tom Fahey ² and Dr. Gráinne Cousins¹

Affiliations:

¹School of Pharmacy, Royal College of Surgeons, 1st Floor Ardilaun House Block B, 111 St Stephen’s Green, Dublin 2, Ireland

²HRB Centre for Primary Care Research, Division of Population Health Science, Royal College of Surgeons in Ireland, 123 St. Stephens Green, Dublin 2, Ireland

Corresponding author: Ms. Alice Holton MPSI, RCSI School of Pharmacy, Block B Ardilaun House, Dublin 2. Email: aliceholton@rcsi.ie

Abstract:

Objectives: Older adults are particularly vulnerable to adverse effects from concurrent alcohol and medication use. However, there is limited evidence regarding the prevalence of these adverse outcomes among older adults and there is a lack of consensus regarding what constitutes an alcohol-interactive medicine. The objective of this study was to develop an explicit list of potentially serious alcohol-medication interactions for use in older adults.

Design: Following a systematic review, review of drug compendia and clinical guidance documents, a two-round Delphi consensus method was conducted.

Setting: Ireland and the United Kingdom (UK), primary care and hospital setting.

Participants: The Project Steering Group developed a list of potentially serious alcohol-medication interactions. The Delphi panel consisted of 19 healthcare professionals (general practitioners (GPs), geriatricians, hospital and community pharmacists, clinical pharmacologists and pharmacists and physicians specialising in substance misuse).

Results: An inventory of 52 potentially serious alcohol-medication interactions was developed by the Project Steering Group. British National Formulary (BNF) black dot warnings (n=8) were included in the final criteria as they represent 'potentially serious' interactions. The remaining 44 criteria underwent a two-round Delphi process. In the first round, thirteen criteria were accepted into the POSAMINO criteria. Consensus was not reached on the remaining 31 criteria; nine were removed, and eight additional criteria were included following a review of panellist comments. The remaining 30 criteria went to round-two, with 17 criteria reaching consensus, providing a final list of 38 potentially serious alcohol-medication interactions; central nervous system (n=15), cardiovascular system (n=9), endocrine system (n=5), musculoskeletal system (n=3), infections (n=3), malignant disease & immunosuppression (n=2) and respiratory system (n=1).

Conclusions: POSAMINO is the first set of explicit potentially serious alcohol-medication interactions for use in older adults. These criteria will facilitate the risk stratification of older adults at the point of prescribing.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Strengths and limitations of this study:

- The potentially serious alcohol-medication interactions for use in older adults (POSAMINO) criteria were developed in a robust fashion, using a two-step process involving a systematic review and two-round Delphi process.
- The Delphi consensus technique utilised in this study is flexible and enabled communication from a diverse group of healthcare professionals from both the United Kingdom and Ireland.
- The final criteria will need to be further validated prospectively to quantify the magnitude of risk posed for each criterion for adverse outcomes in older adults.
- Once validated, these criteria have the potential to inform medical decision making and enable healthcare professionals to risk stratify older adults at the point of prescribing and prioritise alcohol screening and brief interventions in high-risk groups.

Background

Population demographics are changing globally, with the proportion and age of the older population continuing to increase. (1) While alcohol consumption tends to decline in older age, recent evidence suggests that drinking occasions tend to become more frequent among older adults. (2) There is also evidence of a cohort effect with successive birth cohorts reporting an increase in alcohol consumption among all age groups, including older adults. (3) Ageing is associated with a variety of physiological changes, which may place older adults at an increased risk of alcohol-related health problems. (4, 5) In fact, older adults experience a disproportionate burden of alcohol related-harm; in England between 2009 and 2010, adults aged ≥ 65 years accounted for approximately 44% (461,400) of alcohol-related hospital admissions yet comprised of only 17% of the population. (5) Alcohol-related deaths were also highest among those aged 55 to 74 years.(5)

Polypharmacy is also increasing in older adults. (6) For example, approximately 60% of Irish adults aged ≥ 65 years reported taking five or more medicines in 2012. (6) Certain medicines have the potential to interact with alcohol increasing the risk of medical complications such as hypoglycaemia, hypotension, sedation, gastrointestinal bleeds and liver damage. (7, 8) Alcohol interactive (AI) medicines may interact with alcohol by altering the metabolism (pharmacokinetic) or effects (pharmacodynamic) of alcohol and/or the medication. (8) Interactions may occur with any alcohol or follow a dose response, with the risk or severity of an interaction increasing with increasing levels of alcohol consumption. (7)

While a number of studies have investigated the concurrent use of alcohol and medicines with potential to interact with alcohol among older adults, (9-28) there is a lack of consensus regarding what constitutes an AI medication across studies. Several studies focussed on a wide range of medications, using different drug compendia to identify medications as alcohol interactive, thus leading to a lack of consistency in the inclusion of medicines as alcohol interactive. (9-12, 16, 19, 21-23) Other studies focussed on psychotropic medications alone. (13, 15, 17, 18, 20, 24, 26, 28)

While there appears to be a high propensity for alcohol-medication interactions among community dwelling older adults, with between one-in-five to one-in-three older adults potentially susceptible to alcohol-medication interactions, (11, 12, 23) no study to date has examined longitudinal associations of concurrent use

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

with adverse outcomes. An evidence based list of medications which have a significant risk of harm to older patients when combined with alcohol has potential in a clinical setting, once validated, allowing for the identification of older adults whose alcohol consumption places them at increased risk and who would benefit from a preventative intervention. Therefore, the aim of this study is to derive the first set of explicit potentially serious alcohol-medication interactions in older adults.

For peer review only

Methods

Study design

A Delphi consensus technique was used to develop the list of potentially serious alcohol-medication interactions in older adults. The Delphi method allows a consensus opinion to be reached amongst a panel of experts through an iterative process of questionnaires. (29) Ethics approval for this study was obtained from the Royal College of Surgeons in Ireland (Reference Number REC1097). A Project Steering Group comprising of academic and clinical pharmacists, a general practitioner and an epidemiologist (authors) from the RCSI School of Pharmacy and HRB Primary Care Centre was formed to develop the initial list of potentially serious alcohol-medication interactions and to later oversee the Delphi consensus study.

Compilation of initial list of potentially serious alcohol-medication interactions

Following a comprehensive systematic search using MEDLINE (PubMed), Embase, Scopus and Web of Science databases (30), an extensive list of medications with potential to interact with alcohol was identified by the Project Steering Group. A combination of the following keywords and MeSH terms were used: “ethanol”, “alcohol”, “drug interactions”, “drug alcohol interaction” and “aged”. This search was supplemented by a search in Google Scholar and by hand searching references of retrieved articles. The search was restricted to English language articles and articles published since January 1990. Furthermore, the British National Formulary (BNF), Stockley’s Drug Interactions and Martindale Complete Drug Reference drug compendia were also searched. (31-33) Additional documents such as clinical guidance documents (34) and previous reviews (7, 8) were also accessed. Information extracted included: medication name/class, potential adverse outcome(s), whether an interaction is likely to occur with any alcohol consumption or with heavy consumption using national low risk drinking guidelines (35, 36) and, if reported, evidence supporting the interaction. The list of medications was organised according to the BNF physiological classification system. There was considerable heterogeneity across reference sources in terms of identifying medications as having potential to interact with alcohol, with no age specific information for interactions. Furthermore, there were inconsistencies in relation to the quantity of concurrent alcohol consumption that should be avoided. For

example, the Martindale mentioned that alcohol combined with NSAIDs may increase the risk of gastrointestinal bleeds with no mention of alcohol consumption patterns, Stockley's described that both NSAIDs and excessive alcohol use carry the risk of gastrointestinal adverse effects, while NSAIDs and alcohol were not included as an interaction in the BNF. For the next step, the Project Steering Group reviewed and assessed each medication or drug class. Using the definition of "necessary to avoid" previously described by Dreischulte et al. (37), it was considered necessary to avoid concurrent alcohol with a specific medication if, in an average older adult concurrent use of alcohol with this medication would be considered unsafe because (a) there is sufficient evidence that the patient is likely to be harmed, and (b) the likely harm to the patient is large enough to be clinically significant.

Drugs were excluded for the following reasons: their interaction with alcohol was not considered serious i.e. unlikely to cause significant harm to the patient (for example SSRIs and alcohol) (33); the medication is not licensed for use in older adults; the medication is only administered in a hospital environment (for example Trabectedin), interaction with alcohol may only occur during alcohol withdrawal or has been withdrawn from the market in the UK or Ireland. Following these exclusions a truncated list was further reviewed by the Project Steering Group by a consensus discussion. In this process the Project Steering Group agreed, that BNF black dot warnings would be included in the final list. BNF black dots refer to potentially serious drug-alcohol interactions where concurrent use should be avoided or only undertaken with caution and appropriate management. In the paper version of the BNF, these potentially serious drug-alcohol interactions are highlighted to prescribers by flagging them with a 'black dot' (31). Furthermore, the amount of alcohol to be avoided with each medication class was defined through group discussion, based on both the evidence available and the members' own clinical experience. Additionally, it was agreed that the following adverse outcomes associated with concurrent use were defined using clinical guidance documents and reviews to ensure clarity among all participants: orthostatic hypotension (38), hypoglycaemia (39) and lactic acidosis (40).

Selection of the Delphi panel

A total of 51 experts from the Republic of Ireland and United Kingdom, were invited (via e-mail or letter) to participate as part of the Delphi consensus panel. The experts invited to participate on the Delphi consensus panel were peer recognized by the Project Steering Group or nominated by other panel members on the basis of their clinical experience or knowledge of alcohol-medication interactions or care of older persons.

Reasons for non-participation were not required; however in some instances they were provided, they included time commitments and lack of clinical knowledge of alcohol-medication interactions. In total, 19 participants (37%) agreed to participate and written consent was received from all participants before the study. The panel consisted of general practitioners (GPs) (n=5), geriatricians (n=3), hospital pharmacists with expertise in care of older adults (n=3), community pharmacists (n=3), clinical pharmacologists (n=2), clinical pharmacists specialising in substance misuse (n=2) and a physician specialising in substance misuse (n=1). Panel members were not provided with compensation for participation.

The Delphi validation technique and process

An online questionnaire was piloted by two pharmacists and two GPs, to identify any potential problems and to approximate the completion time for the survey. Following amendments, the 19 participants were sent a link to the online POSAMINO questionnaire (via SurveyGizmo®) in March 2016. Participants were given four weeks to complete the online questionnaire and all participants were sent a reminder email after 2 weeks to encourage participation. The questionnaire was both anonymous and confidential.

The quantity of alcohol per standard drink (SD) (10 grammes alcohol in Ireland) or unit (8 grammes alcohol in the United Kingdom), was defined at the beginning of the survey. As some interactions are listed with any alcohol consumption and others with 'heavy' consumption, we provided definitions of both at the beginning of the questionnaire. Any alcohol consumption was defined as ≥ 1 SD of alcohol in Ireland or ≥ 1 unit of alcohol in the UK, with heavy consumption defined according to Irish and UK National low risk drinking guidelines. (35, 36) The panel were instructed to evaluate each potential interaction listed according to the "necessary to avoid" framework described above. (37) Medications were categorised according to the BNF

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

physiological classification system. Each statement was presented in the same format stating: It is necessary to avoid [quantity of alcohol] (any or heavy) with [medications/ drug class], followed by a brief rationale for the statement. For example: *It is necessary to avoid: Any alcohol consumption with first generation anti-histamines (for example promethazine). (Rationale: Concurrent alcohol consumption with first generation anti-histamines can increase the risk of sedation).*

Participants were asked to rate their agreement with each statement, using a 5-point Likert scale (1-strongly disagree to 5-strongly agree) along with any additional comments, or suggestions for additional medicines to be included.

After each round, the median and interquartile ranges (IQR) were calculated for each statement. Consistent with previous Delphi consensus studies, the consensus level required for a statement to be retained was defined *a priori* as a median of 4 or 5 with a lower quartile value of ≥ 4 . (41) If a statement had an upper quartile value of ≤ 2 , this indicated there was general disagreement with the statement between panel members, and the statement was rejected. If group consensus was not reached, the criteria were reviewed by the Project Steering Group and were removed based on comments or revised and included in the second questionnaire. In the second questionnaire, panellists were provided with links to the most recent evidence relating to each of the alcohol-medication interactions, to help facilitate their decisions due to uncertainty in round 1. As with round one, the median response and IQR were calculated and evaluated by the Project Steering Group using the same thresholds to determine consensus between the panel members. If consensus was not reached following the second round, the statement was rejected. Statistical analysis was performed using STATA version 13 and Microsoft Excel 2010.

Results

Following an initial review of the literature, Anatomical Therapeutic Chemical (ATC) codes for a total of 364 AI medicines were extracted by the Project Steering Group (**Supplementary File 1**). Medicines were classified according to drug classes and organised according to BNF physiological classification systems. From this list, a total of 63 statements were initially compiled based on the ‘necessary to avoid’ framework. (37) Following group discussions, 11 statements were removed/updated or merged together, based on the steering groups clinical knowledge or experience with medicines. For example, the steering group decided to no longer classify benzodiazepines according to the duration of action, as both long acting and short acting have the potential to interact with alcohol. Furthermore, a total of eight BNF black dot warnings were included in the final criteria as they represent potentially serious interactions.

All 19 panel members completed the round one questionnaire, in which forty-four statements were presented. Consensus was reached on 13 statements, with no statements rejected (**Table 1**). Thirty one statements were reviewed by the steering group, with nine statements removed based on comments from the panel, if the interaction was not relevant to older adults or the interaction was not of clinical significance for example: heavy alcohol consumption with vitamin A preparations. Furthermore, eight new statements were included in round two based on comments from panellists at the first round.

A total of 18 out of the 19 recruited participants completed the round two questionnaire. Of the 30 statements included in the second questionnaire, consensus was reached for 17 statements. The remaining 13 statements were rejected as no consensus was reached. In total, consensus was reached for 30 potentially serious drug alcohol interactions in older adults, with the inclusion of BNF black dot warnings the final list was 38 statements.

INSERT TABLE 1 HERE

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

This final 38 item POSAMINO criteria (**Table 2**) were organised over the following physiological systems:
Central nervous system (n=15), cardiovascular system (n=9), endocrine system (n=5), musculoskeletal system
(n=3), infections (n=3), malignant disease and immunosuppression (n=2) and respiratory system (n=1).

INSERT TABLE 2 HERE

For peer review only

Discussion

Principle findings in relation to previous studies

While older adults experience a disproportionate burden of alcohol related harm, (5) research suggests that healthcare professionals are less likely to discuss alcohol consumption with older adults. (42, 43) Flagging older adults at the point of prescribing an alcohol-interactive medication may facilitate targeted screening and interventions to help reduce harm. Despite the high propensity for alcohol-medication interactions among community dwelling older adults, our recent systematic review has highlighted that there is still a lack of consensus regarding what constitutes an AI medication.(30)

This study reports the development of a set of 38 explicit criteria, POSAMINO, for identifying older adults at risk of potentially serious alcohol-medication interactions. The final POSAMINO criteria consist of seven different drug classes, with central nervous system (CNS) medicines representing 40% of the criteria. Estimates from our previous study indicate that approximately one in five older Irish adults are exposed to CNS agents, with over half of patients using CNS agents also reporting concurrent alcohol consumption. (12) Nine of the final 38 criteria concern cardiovascular agents, another widely used drug class among aged adults. (12) There is also a high-risk of adverse effects associated with these agents, with anti-platelets, diuretics and anticoagulants identified as the most common drug classes involved in preventable drug-related admissions, in a previous study. (44) Analgesics such as non-steroidal anti-inflammatory drugs (NSAIDs), opioids, paracetamol and gabapentin (used for neuropathic pain) were also included in the final POSAMINO criteria. (44) The prevalence of chronic pain increases with age and older adults may consume alcohol to help cope and manage their pain. (45) This behaviour may also predispose older adults to adverse outcomes associated with concurrent use with analgesics (for example increased sedation, falls and gastrointestinal bleeds).

The development of the POSAMINO criteria is important for several reasons. In the absence of an explicit list of alcohol-interactive medications, multiple drug reference sources have been used in previous studies leading to a lack of consistency in the inclusion of medications across studies, (11, 12, 19, 22, 23) which may result in biased estimates of potential risk among older adults. Furthermore, some of these interactions may be theoretical with trivial clinical significance. Therefore, this study developed a consensus-based set of explicit

criteria for potentially serious alcohol-medication interactions in older adults rather than an exhaustive list of medications with potential to interact with alcohol. Consistent with existing literature, the POSAMINO criteria classify CNS agents as AI medications with the potential to cause serious harm to older adults (9-13, 15-24, 26, 28). However, some of the previous studies focussed on sedatives/hypnotics only (13, 26, 28). We identified a number of additional CNS agents, such as anti-Parkinson's drugs (for example: Pramipexole, Apomorphine and Levodopa), which may be overlooked if we focus solely on sedatives or hypnotics.

Furthermore, the POSAMINO criteria also includes a wide range of other drug classes, such as cardiovascular, respiratory system, musculoskeletal, malignant disease, infections or endocrine agents which were not all considered among previous studies (9-11, 16, 19, 21, 23). It is also important to note that there has been little emphasis on the adverse outcomes or severity of these potential interactions.(22, 30) The development of the POSAMINO criteria may help identify older adults at risk of potentially serious alcohol-medicine interactions in the future.

Clinical implications and future research

This study adds to a growing body of research investigating the concurrent use of alcohol and alcohol interactive medicines in older adults. Undoubtedly, adverse drug events (ADEs) represent a major burden on healthcare, with ADEs detected in 26.3% of patients aged ≥ 65 years admitted to an Irish hospital with acute illnesses. (46) The increasing use of multiple medications, (6) combined with an increased frequency of alcohol consumption (2) and age-related physiological changes, may predispose older adults to these adverse events. The significant burden of alcohol-related harm and mortality among older adults, (5) indicates a pressing need for future interventions to minimise risk. However, prior to informing clinical or public health initiatives, the final criteria developed from this study, will require further validation to prospectively quantify the magnitude of risk posed for each criterion for adverse outcomes in older adults. In particular, the quantity of alcohol specified in each criterion will need to be further evaluated in prospective studies. Once validated, these criteria have the potential to inform medical decision making and enable healthcare professionals to risk stratify older adults at the point of prescribing and prioritise alcohol screening and brief interventions in high-

1
2
3 risk groups. This is essential, as older people have previously reported little knowledge about risks associated
4 with the concurrent use of alcohol and medications. (47)

5 6 7 8 *Strengths and Limitations*

9
10 The criteria were developed in a robust fashion, using a two-step process involving a systematic review and
11 two-round Delphi process. The Delphi technique is flexible and for this study enabled communication from a
12 group of nineteen healthcare professionals from both the United Kingdom and Ireland. Eighteen of the 19
13 participants completed the two rounds, with participants providing numerous comments and suggestions in
14 both rounds. Similar to previous studies, participants remained anonymous and were not provided with
15 feedback following the first round in order to remove the risk of potential bias. (41)

16
17 Inevitably, there were limitations to this Delphi study. With all Delphi studies, participants' judgement may be
18 subjective. In order to reduce this potential bias, a diverse group of healthcare professionals with expertise or
19 interest in the care of older adults were selected. With explicit criteria, there is also a need for regular updates
20 and revision due to the availability of more up to date information after development. (48) The developed
21 POSAMINO criteria do not apply to older adults diagnosed with an alcohol use disorder, as chronic heavy
22 alcohol consumption can substantially increase the activity of the cytochrome P450 metabolising enzyme
23 CYP2E1. (8) Finally, older adults may also experience chronic illnesses which affect the metabolism of both
24 alcohol and medications. As a result, it is important that healthcare professionals also consider these comorbid
25 diseases when assessing the risk for potential adverse outcomes. (49)

26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 *Conclusions*

44
45 Using a systematic review and a two-round Delphi consensus method, we have developed the first set of
46 explicit potentially serious alcohol-medication interactions (POSAMINO) for use in older adults. Following
47 future validation studies, these criteria may allow for the risk stratification of older adults at the point of
48 prescribing, and prioritise alcohol screening and brief alcohol interventions in high-risk groups.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Declarations:

Authors’ contributions: AH, PG, CR, TF & GC (the project steering group) conceived and designed this study. AH & GC conducted initial literature search. AH communicated with Delphi participants. AH analysed the data after each round. All Project Steering Group members were involved in interpretation of the data. AH & GC drafted the manuscript. All co-authors revised the manuscript and have given the approval for publication.

Competing interests: The authors declare that they have no competing interests.

Ethics approval: Ethical approval for this study was approved by the Royal College of Surgeons in Ireland (RCSI) ethics committee (REC 1097).

Acknowledgements: We would like to thank all Delphi participants who contributed to the development of the POSAMINO criteria.

Funding: This study was funded by the Royal College of Surgeons in Ireland (RCSI), School of Pharmacy, Dublin 2, Ireland.

Data sharing statement: Additional data is available by request from the corresponding author.

References

1. United Nations, Department of Economic and Social Affairs PD. World Population Prospects: The 2015 Revision, Key Findings and Advance Tables. New York: 2015. Available from: https://esa.un.org/unpd/wpp/Publications/Files/Key_Findings_WPP_2015.pdf
2. Britton A, Ben-Shlomo Y, Benzeval M, Kuh D, Bell S. Life course trajectories of alcohol consumption in the United Kingdom using longitudinal data from nine cohort studies. *BMC Med*. 2015;13:47.
3. Meng Y, Holmes J, Hill-McManus D, Brennan A, Meier PS. Trend analysis and modelling of gender-specific age, period and birth cohort effects on alcohol abstinence and consumption level for drinkers in Great Britain using the General Lifestyle Survey 1984–2009. *Addiction*. 2014;109.
4. Huang AR, Mallet L, Rochefort CM, Eguale T, Buckeridge DL, Tamblyn R. Medication-related falls in the elderly: causative factors and preventive strategies. *Drugs Aging*. 2012;29(5):359-76.
5. Wadd S, Papadopoulos C. Drinking behaviour and alcohol-related harm amongst older adults: analysis of existing UK datasets. *BMC Res Notes*. 2014;7:741.
6. Moriarty F, Hardy C, Bennett K, Smith SM, Fahey T. Trends and interaction of polypharmacy and potentially inappropriate prescribing in primary care over 15 years in Ireland: a repeated cross-sectional study. *BMJ open*. 2015;5.
7. Moore AA, Whiteman EJ, Ward KT. Risks of combined alcohol/medication use in older adults. *Am J Geriatr Pharmacother*. 2007;5.
8. Weathermon R, Crabb DW. Alcohol and medication interactions. *Alcohol research & health : the journal of the National Institute on Alcohol Abuse and Alcoholism*. 1999;23(1):40-54.
9. Adams WL. Potential for adverse drug-alcohol interactions among retirement community residents. *J Am Geriatr Soc*. 1995;43.
10. Aira M, Hartikainen S, Sulkava R. Community prevalence of alcohol use and concomitant use of medication - A source of possible risk in the elderly aged 75 and older? *Int J Geriatr Psychiatry*. 2005;20(7):680-5.
11. Breslow RA, Dong C, White A. Prevalence of alcohol-interactive prescription medication use among current drinkers: United States, 1999 to 2010. *Alcohol Clin Exp Res*. 2015;39(2):371-9.
12. Cousins G, Galvin R, Flood M, Kennedy M-C, Motterlini N, Henman MC, et al. Potential for alcohol and drug interactions in older adults: evidence from the Irish longitudinal study on ageing. *BMC Geriatr*. 2014;14.
13. Del Rio MC, Prada C, Alvarez FJ. Do Spanish patients drink alcohol while undergoing treatment with benzodiazepines? *Alcohol*. 2002;26(1):31-4.

14. Del Río MC, Prada C, Alvarez FJ. The use of medication and alcohol among the Spanish population. *Br J Clin Pharmacol*. 1996;41(3):253-5.

15. Du Y, Scheidt-Nave C, Knopf H. Use of psychotropic drugs and alcohol among non-institutionalised elderly adults in Germany. *Pharmacopsychiatry*. 2008;41(6):242-51.

16. Forster LE, Pollock R, Stoller EP. Alcohol use and potential risk for alcohol-related adverse drug reactions among community-based elderly. *J Community Health*. 1993;18(4):225-39.

17. Ilomaki J, Korhonen MJ, Enlund H, Hartzema AG, Kauhanen J. Risk drinking behavior among psychotropic drug users in an aging Finnish population: the FinDrink study. *Alcohol*. 2008;42(4):261-7.

18. Ilomaki J, Gnjdic D, Hilmer SN, Le Couteur DG, Naganathan V, Cumming RG, et al. Psychotropic drug use and alcohol drinking in community-dwelling older Australian men: the CHAMP study. *Drug and alcohol review*. 2013;32(2):218-22.

19. Immonen S, Pitkälä K. The prevalence of potential alcohol-drug interactions in older adults. *Eur Geriatr Med*. 2012;3:S134.

20. John U, Baumeister SE, Volzke H, Meyer C, Ulbricht S, Alte D. Sedative, hypnotic, anxiolytic and opioid medicament use and its co-occurrence with tobacco smoking and alcohol risk drinking in a community sample. *BMC Public Health*. 2007;7:337.

21. Onder G, Landi F, Della Vedova C, Atkinson H, Pedone C, Cesari M, et al. Moderate alcohol consumption and adverse drug reactions among older adults. *Pharmacoepidemiol Drug Saf*. 2002;11(5):385-92.

22. Pringle KE, Ahern FM, Heller DA, Gold CH, Brown TV. Potential for alcohol and prescription drug interactions in older people. *J Am Geriatr Soc*. 2005;53.

23. Qato DM, Manzoor BS, Lee TA. Drug-alcohol interactions in older US adults. *J Am Geriatr Soc*. 2015;63.

24. Sheahan SL, Coons SJ, Robbins CA, Martin SS, Hendricks J, Latimer M. Psychoactive Medication, Alcohol-Use, and Falls among Older Adults. *J Behav Med*. 1995;18(2):127-40.

25. Swift W, Stollznow N, Pirotta M. The use of alcohol and medicines among Australian adults. *Aust N Z J Public Health*. 2007;31.

26. Veldhuizen S, Wade TJ, Cairney J. Alcohol consumption among Canadians taking benzodiazepines and related drugs. *Pharmacoepidemiol Drug Saf*. 2009;18.

27. Wong H, Heuberger R, Logomarsino J, Hewlings S. Associations between alcohol use, polypharmacy and falls in older adults. *Nurs Older People*. 2016;28(1):30-6.

28. Lagnaoui R, Moore N, Dartigues JF, Fourrier A, Begaud B. Benzodiazepine use and wine consumption in the French elderly. *Br J Clin Pharmacol*. 2001;52(4):455-6.

29. Hsu C-C, Sandford BA. The Delphi technique: making sense of consensus. Practical assessment, research & evaluation. 12(10):1-8.
30. Holton AE, Gallagher P, Fahey T, Cousins G. Concurrent use of alcohol interactive medications and alcohol in older adults: a systematic review of prevalence and associated adverse outcomes. BMC Geriatr. 2017;17(1):148.
31. Joint Formulary Committee (2015) The British National Formulary (BNF). 69th Ed. London- British Medical Association and Royal Pharmaceutical Society of Great Britain
32. Sweetman S. Martindale: The Complete Drug Reference London: Pharmaceutical Press; [cited 2015]. Available from: <https://www.medicinescomplete.com.proxy.library.rcsi.ie/mc/martindale/current/index.htm>.
33. Stockley's Drug Interactions [online] [Internet]. Pharmaceutical Press. Available from: <https://www.medicinescomplete.com/mc/stockley/current/>.
34. National Institute on Alcohol Abuse and Alcoholism (NIAAA). Harmful Interactions mixing alcohol with medicines. 2014. Available from: https://pubs.niaaa.nih.gov/publications/medicine/Harmful_Interactions.pdf
35. Hope. A Standard Drink in Ireland: What strength? Health Service Executive- Alcohol Implementation Group 2009
36. UK Chief Medical Officers' Alcohol Guidelines Review Summary of the proposed new guidelines. Department of Health January 2016 Available from: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/489795/summary.pdf
37. Dreischulte T, Grant AM, McCowan C, McAnaw JJ, Guthrie B. Quality and safety of medication use in primary care: consensus validation of a new set of explicit medication assessment criteria and prioritisation of topics for improvement. BMC Clin Pharmacol. 2012;12:5.
38. Lanier JB, Mote MB, Clay EC. Evaluation and Management of Orthostatic Hypotension. Am Fam Physician. 2011;84(5):527-36.
39. European Medicines Agency (EMA). Guideline on clinical investigation of medicinal products in the treatment or prevention of diabetes mellitus Committee for Medicinal Products for Human Use, 2012 Available from: http://www.ema.europa.eu/docs/en_GB/document_library/Scientific_guideline/2012/06/WC500129256.pdf
40. Salpeter SR, Greyber E, Pasternak GA, Salpeter EE. Risk of fatal and nonfatal lactic acidosis with metformin use in type 2 diabetes mellitus: Systematic review and meta-analysis. Arch Intern Med. 2003;163(21):2594-602.
41. Cooper JA, Ryan C, Smith SM, Wallace E, Bennett K, Cahir C, et al. The development of the PROMPT (PRescribing Optimally in Middle-aged People's Treatments) criteria. BMC Health Serv Res. 2014;14:484.

42. Duru OK, Xu H, Tseng C-H, Mirkin M, Ang A, Tallen L, et al. Correlates of Alcohol-Related Discussions between Older Adults and Their Physicians. *J Am Geriatr Soc.* 2010;58(12):2369-74.

43. Kuerbis A, Sacco P, Blazer DG, Moore AA. Substance Abuse Among Older Adults. *Clin Geriatr Med.* 2014;30(3):629-54.

44. Howard RL, Avery AJ, Slavenburg S, Royal S, Pipe G, Lucassen P, et al. Which drugs cause preventable admissions to hospital? A systematic review. *Br J Clin Pharmacol.* 2007;63(2):136-47.

45. Ahangari A, Stewart Williams J, Myleus A. Pain and alcohol consumption among older adults: findings from the World Health Organization Study on global AGEing and adult health, Wave 1. *Trop Med Int Health.* 2016;21(10):1282-92.

46. Hamilton H, Gallagher P, Ryan C, Byrne S, O'Mahony D. Potentially inappropriate medications defined by stopp criteria and the risk of adverse drug events in older hospitalized patients. *Arch Intern Med.* 2011;171(11):1013-9.

47. Zanjani F, Hoogland AI, Downer BG. Alcohol and prescription drug safety in older adults. *Drug, healthcare and patient safety.* 2013;5:13-27.

48. O'Mahony D, O'Sullivan D, Byrne S, O'Connor MN, Ryan C, Gallagher P. STOPP/START criteria for potentially inappropriate prescribing in older people: version 2. *Age Ageing.* 2015;44(2):213-8.

49. Slattum PW, Hassan OE. Medications, Alcohol, and Aging. In: Kuerbis A, Moore AA, Sacco P, Zanjani F, editors. *Alcohol and Aging: Clinical and Public Health Perspectives.* Cham: Springer International Publishing; 2016. p. 117-29.

Table 1: Results from the 2-round Delphi process

	Potentially serious interactions as listed in the BNF	Round 1					Round 2					Final Criteria
		Total	Accept	Revise	Reject	Removal	New statements included	Total	Accept	Revise	Reject	
Physiological System												
Cardiovascular System	1	14	1	13	0	3	2	12	7	0	5	9
Respiratory System	0	1	1	0	0	0	0	0	0	0	0	1
Central Nervous System	4	12	6	6	0	1	5	10	5	0	5	15
Infections	2	5	0	5	0	3	0	2	1	0	1	3
Endocrine System	0	5	3	2	0	0	0	2	2	0	0	5
Obstetrics, Gynaecology, and Urinary Tract Disorders	0	2	0	2	0	1	1	2	0	0	2	0
Malignant Disease and Immunosuppression	1	1	0	1	0	0	0	1	1	0	0	2
Nutrition and Blood	0	1	0	1	0	1	0	0	0	0	0	0
Musculoskeletal System	0	3	2	1	0	0	0	1	1	0	0	3
Total	8	44	13	31	0	9	8	30	17	0	13	38

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 2: Final POSAMINO Criteria (n=38)

Cardiovascular System

- 1. **Heavy alcohol consumption** with multiple anti-hypertensive combinations
Rationale: Concurrent use of alcohol consumption and anti-hypertensives may increase the risk of orthostatic hypotension
- 2. **Heavy alcohol consumption** with warfarin (& phenindione)
Rationale: Heavy episodic alcohol consumption is associated with an increased risk of major bleeds
- 3. **Heavy alcohol consumption** with regular use of low dose aspirin (75mg)
Rationale: Heavy alcohol consumption combined with aspirin may cause a small increase in gastrointestinal blood loss
- 4. **Heavy alcohol consumption** with both regular and as required nitrates (e.g. glyceryl trinitrate, isosorbide dinitrate and isosorbide mononitrate)
Rationale: The combined haemodynamic effects of alcohol and nitrates, increases the risk of exaggerated hypotension.
- 5. **Heavy alcohol consumption** with the vasodilatory medication nicorandil
Rationale: The combined haemodynamic effects of alcohol & nicorandil; increases the risk of exaggerated hypotension.
- 6. **Heavy alcohol consumption** with the combined use of both nitrates and vasodilator medication (e.g. nicorandil)
Rationale: The combined haemodynamic effects of alcohol and nitrates, increases the risk of exaggerated hypotension.
- 7. **Heavy alcohol consumption** with diuretics (e.g. loop diuretics (furosemide), thiazide diuretics (bendroflumethiazide) & potassium sparing diuretics (amiloride))
Rationale: The concurrent use of alcohol and anti-hypertensives may increase the risk of orthostatic hypotension.
- 8. **Heavy alcohol consumption** with alpha blockers (e.g. terazosin)
Rationale: The concurrent use of alcohol and anti-hypertensives may increase the risk of orthostatic hypotension.
- 9. **Heavy alcohol consumption** with centrally acting anti-hypertensives (e.g. clonidine or methyldopa)
Rationale: Alcohol consumption combined with centrally acting anti-hypertensives may increase the risk of sedation.

Respiratory System

- 1. **Any alcohol consumption** with first generation anti-histamines (e.g. promethazine)
Rationale: Concurrent alcohol consumption with first generation anti-histamines can increase the risk of sedation

Central Nervous System

- 1. **Heavy alcohol consumption** with benzodiazepines (e.g.diazepam) and benzodiazepine related medications (e.g.zopiclone)
Rationale: Alcohol consumption combined with benzodiazepines and benzodiazepine related medications may enhance CNS depressant effects
- 2. **Heavy alcohol consumption** combined with opioids
Rationale: Alcohol consumption combined with opioids may enhance CNS depressant effects of alcohol
- 3. **Heavy alcohol consumption** with duloxetine
Rationale: Heavy alcohol consumption combined with duloxetine may increase the risk of hepatotoxicity
- 4. **Heavy alcohol consumption** with all anti-psychotics
Rationale: Alcohol consumption combined with antipsychotics may increase the risk of sedation
- 5. **Any alcohol consumption** with barbiturates
Rationale: Alcohol consumption combined with barbiturates may increase the risk of sedation.
- 6. **Heavy alcohol consumption** with anti-epileptic drugs (AEDs)
Rationale: Heavy alcohol consumption can increase the risk of seizures and sedation in patients taking AEDs.
- 7. **Any alcohol consumption** with tricyclic anti-depressants (TCAs)
Rationale: Alcohol consumption combined with antidepressants may enhance the CNS depressant effects of alcohol
- 8. **Any alcohol consumption** with tetracyclic antidepressants
Rationale: Alcohol consumption combined with antidepressants may enhance the CNS depressant effects of alcohol
- 9. **Any alcohol consumption** with mirtazapine
Rationale: Alcohol consumption combined with antidepressants may enhance the CNS depressant effects of alcohol
- 10. **Any alcohol consumption** with monoamine oxidase inhibitors (MAOIs)
Rationale: A potentially life-threatening hypertensive reaction can develop in patients taking non-selective MAOIs who consume drinks rich in tyramine (e.g. wines, beers and non-alcoholic beers)
- 11. **Heavy alcohol consumption** with long term regular paracetamol use (e.g. 1g four times a day)
Rationale: Heavy alcohol consumption may increase the risk of paracetamol hepatotoxicity especially if alcohol intake is abruptly stopped.
- 12. **Heavy alcohol consumption** with gabapentin (when used for neuropathic pain)
Rationale: Alcohol combined with medications for neuropathic pain may increase the risk of sedation
- 13. **Heavy alcohol consumption** with pramipexole or amantadine
Rationale: Alcohol may increase the risk of sedation
- 14. **Heavy alcohol consumption** with apomorphine
Rationale: Alcohol combined with apomorphine may increase the risk of orthostatic hypotension

15. Heavy alcohol consumption with levodopa (alone or in combination with carbidopa)

Rationale: Alcohol combined with levodopa (alone or in combination with carbidopa) may increase the risk of orthostatic hypotension.

Endocrine

1. Heavy alcohol consumption with insulin

Rationale: Alcohol consumption may enhance the hypoglycaemic effect of insulin.

2. Heavy alcohol consumption with metformin

Rationale: Heavy alcohol consumption combined with metformin may increase the risk of lactic acidosis

3. Heavy alcohol consumption with sulphonylureas

Rationale: Alcohol consumption can enhance the hypoglycaemic effects of antidiabetics

4. Heavy alcohol consumption with meglitinides (e.g. nateglinide)

Rationale: Alcohol consumption can enhance the hypoglycaemic effects of antidiabetics

5. Heavy alcohol consumption with thiazolidinediones (e.g. pioglitazone)

Rationale: Alcohol consumption can enhance the hypoglycaemic effects of antidiabetics

Musculoskeletal and joint diseases

1. Heavy alcohol consumption with any nonsteroidal anti-inflammatory drugs NSAIDs (including COX-2 inhibitors)

Rationale: Heavy alcohol consumption and NSAID use carry an increased risk of gastrointestinal bleeds

2. Heavy alcohol consumption combined with methotrexate or leflunomide

Rationale: Heavy alcohol consumption combined with methotrexate or leflunomide may increase the risk of hepatotoxicity

3. Heavy alcohol consumption with oral muscle relaxants (e.g. baclofen)

Rationale: Concurrent alcohol consumption and muscle relaxants can increase the risk of CNS depression.

Malignant disease and immunosuppression

1. Any alcohol consumption with procarbazine

Rationale: A disulfiram-like reaction can occur when alcohol is given with procarbazine

2. Heavy alcohol consumption with interferon alpha or interferon beta

Rationale: Heavy alcohol consumption combined with interferons may increase the risk of hepatotoxicity and reduce the response to treatment with interferon.

Infections

1. Heavy alcohol consumption with anti-mycobacterial medications such as isoniazid, pyrazinamide, ethionamide and rifampicin (alone or in combination)

Rationale: Alcohol combined with anti-mycobacterial medications increase the risk of hepatotoxicity

2. Any alcohol consumption with cycloserine

Rationale: Alcohol consumption may increase the risk of seizures in patients taking cycloserine.

3. Any alcohol consumption with metronidazole or tinidazole

Rationale: A disulfiram-like reaction can occur when alcohol is given with metronidazole




Figure 1: Flow chart summarizing the steps involved in the development of the POSAMINO criteria

BMJ Open

Consensus Validation of the POSAMINO (POtentially Serious Alcohol Medication INTERactions in Older adults) Criteria

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-017453.R2
Article Type:	Research
Date Submitted by the Author:	06-Sep-2017
Complete List of Authors:	Holton, Alice; School of Pharmacy Gallagher, Paul; Royal College of Surgeons in Ireland, School of Pharmacy Ryan, Cristin; Royal College of Surgeons in Ireland, Fahey, Tom; Royal College of Surgeons in Ireland, Department of General Practice Cousins, Gráinne; Royal College of Surgeons in Ireland, School of Pharmacy
Primary Subject Heading:	Pharmacology and therapeutics
Secondary Subject Heading:	Health services research, Geriatric medicine
Keywords:	CLINICAL PHARMACOLOGY, Adverse events < THERAPEUTICS, alcohol, GERIATRIC MEDICINE

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Consensus Validation of the POSAMINO (POtentially Serious Alcohol Medication INteractions in Older adults) Criteria

Ms. Alice E. Holton ¹, Prof. Paul Gallagher¹, Dr. Cristín Ryan¹, Prof. Tom Fahey ² and Dr. Gráinne Cousins¹

Affiliations:

¹School of Pharmacy, Royal College of Surgeons, 1st Floor Ardilaun House Block B, 111 St Stephen’s Green, Dublin 2, Ireland

²HRB Centre for Primary Care Research, Division of Population Health Science, Royal College of Surgeons in Ireland, 123 St. Stephens Green, Dublin 2, Ireland

Corresponding author: Ms. Alice Holton MPSI, RCSI School of Pharmacy, Block B Ardilaun House, Dublin 2. Email: aliceholton@rcsi.ie

Abstract:

Objectives: Older adults are particularly vulnerable to adverse effects from concurrent alcohol and medication use. However, there is limited evidence regarding the prevalence of these adverse outcomes among older adults and there is a lack of consensus regarding what constitutes an alcohol-interactive medicine. The objective of this study was to develop an explicit list of potentially serious alcohol-medication interactions for use in older adults.

Design: Following a systematic review, review of drug compendia and clinical guidance documents, a two-round Delphi consensus method was conducted.

Setting: Ireland and the United Kingdom (UK), primary care and hospital setting.

Participants: The Project Steering Group developed a list of potentially serious alcohol-medication interactions. The Delphi panel consisted of 19 healthcare professionals (general practitioners (GPs), geriatricians, hospital and community pharmacists, clinical pharmacologists and pharmacists and physicians specialising in substance misuse).

Results: An inventory of 52 potentially serious alcohol-medication interactions was developed by the Project Steering Group. British National Formulary (BNF) black dot warnings (n=8) were included in the final criteria as they represent 'potentially serious' interactions. The remaining 44 criteria underwent a two-round Delphi process. In the first round, thirteen criteria were accepted into the POSAMINO criteria. Consensus was not reached on the remaining 31 criteria; nine were removed, and eight additional criteria were included following a review of panellist comments. The remaining 30 criteria went to round-two, with 17 criteria reaching consensus, providing a final list of 38 potentially serious alcohol-medication interactions; central nervous system (n=15), cardiovascular system (n=9), endocrine system (n=5), musculoskeletal system (n=3), infections (n=3), malignant disease & immunosuppression (n=2) and respiratory system (n=1).

Conclusions: POSAMINO is the first set of explicit potentially serious alcohol-medication interactions for use in older adults. Following future validation studies, these criteria may allow for the risk stratification of older adults at the point of prescribing.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Strengths and limitations of this study:

- The potentially serious alcohol-medication interactions for use in older adults (POSAMINO) criteria were developed in a robust fashion, using a two-step process involving a systematic review and two-round Delphi process.
- The Delphi consensus technique utilised in this study is flexible and enabled communication from a diverse group of healthcare professionals from both the United Kingdom and Ireland.
- The final criteria will need to be further validated prospectively to quantify the magnitude of risk posed for each criterion for adverse outcomes in older adults.
- Once validated, these criteria have the potential to inform medical decision making and enable healthcare professionals to risk stratify older adults at the point of prescribing and prioritise alcohol screening and brief interventions in high-risk groups.

Background

Population demographics are changing globally, with the proportion and age of the older population continuing to increase. (1) While alcohol consumption tends to decline in older age, recent evidence suggests that drinking occasions tend to become more frequent among older adults. (2) There is also evidence of a cohort effect with successive birth cohorts reporting an increase in alcohol consumption among all age groups, including older adults. (3) Ageing is associated with a variety of physiological changes, which may place older adults at an increased risk of alcohol-related health problems. (4, 5) In fact, older adults experience a disproportionate burden of alcohol related-harm; in England between 2009 and 2010, adults aged ≥ 65 years accounted for approximately 44% (461,400) of alcohol-related hospital admissions yet comprised of only 17% of the population. (5) Alcohol-related deaths were also highest among those aged 55 to 74 years.(5)

Polypharmacy is also increasing in older adults. (6) For example, approximately 60% of Irish adults aged ≥ 65 years reported taking five or more medicines in 2012. (6) Certain medicines have the potential to interact with alcohol increasing the risk of medical complications such as hypoglycaemia, hypotension, sedation, gastrointestinal bleeds and liver damage. (7, 8) In one cross sectional study, after adjusting for potential confounders, moderate alcohol consumption was associated with a 24% increased risk in adverse drug reactions (ADRs) among older adults. (21) Alcohol interactive (AI) medicines may interact with alcohol by altering the metabolism (pharmacokinetic) or effects (pharmacodynamic) of alcohol and/or the medication. (8) Interactions may occur with any alcohol or follow a dose response, with the risk or severity of an interaction increasing with increasing levels of alcohol consumption. (7)

While a number of studies have investigated the concurrent use of alcohol and medicines with potential to interact with alcohol among older adults, (9-28) there is a lack of consensus regarding what constitutes an AI medication across studies. Several studies focussed on a wide range of medications, using different drug compendia to identify medications as alcohol interactive, thus leading to a lack of consistency in the inclusion of medicines as alcohol interactive. (9-12, 16, 19, 21-23) Other studies focussed on psychotropic medications alone. (13, 15, 17, 18, 20, 24, 26, 28)

While there appears to be a high propensity for alcohol-medication interactions among community dwelling older adults, with between one-in-five to one-in-three older adults potentially susceptible to alcohol-medication interactions, (11, 12, 23) no study to date has examined longitudinal associations of concurrent use with adverse outcomes. An evidence based list of medications which have a significant risk of harm to older patients when combined with alcohol has potential in a clinical setting, once validated, allowing for the identification of older adults whose alcohol consumption places them at increased risk and who would benefit from a preventative intervention. Therefore, the aim of this study is to derive the first set of explicit potentially serious alcohol-medication interactions in older adults.

Methods

Study design

A Delphi consensus technique was used to develop the list of potentially serious alcohol-medication interactions in older adults. The Delphi method allows a consensus opinion to be reached amongst a panel of experts through an iterative process of questionnaires. (29) Ethics approval for this study was obtained from the Royal College of Surgeons in Ireland (RCSI) (Reference Number REC1097). A Project Steering Group comprising of academic and clinical pharmacists, a general practitioner and an epidemiologist (authors) from the RCSI School of Pharmacy and Health Research Board (HRB) Primary Care Centre was formed to develop the initial list of potentially serious alcohol-medication interactions and to later oversee the Delphi consensus study.

Compilation of initial list of potentially serious alcohol-medication interactions

Following a comprehensive systematic search using MEDLINE (PubMed), Embase, Scopus and Web of Science databases (30), an extensive list of medications with potential to interact with alcohol was identified by the Project Steering Group. A combination of the following keywords and MeSH terms were used: “ethanol”, “alcohol”, “drug interactions”, “drug alcohol interaction” and “aged”. This search was supplemented by a search in Google Scholar and by hand searching references of retrieved articles. The search was restricted to English language articles and articles published since January 1990. Furthermore, the British National Formulary (BNF), Stockley’s Drug Interactions and Martindale Complete Drug Reference drug compendia were also searched. (31-33) Additional documents such as clinical guidance documents (34) and previous reviews (7, 8) were also accessed. Information extracted included: medication name/class, potential adverse outcome(s), whether an interaction is likely to occur with any alcohol consumption or with heavy consumption using national low risk drinking guidelines (35, 36) and, if reported, evidence supporting the interaction. The list of medications was organised according to the BNF physiological classification system. There was considerable heterogeneity across reference sources in terms of identifying medications as having potential to interact with alcohol, with no age specific information for interactions. Furthermore, there were

inconsistencies in relation to the quantity of concurrent alcohol consumption that should be avoided. For example, the Martindale mentioned that alcohol combined with nonsteroidal anti-inflammatory drugs (NSAIDs) may increase the risk of gastrointestinal bleeds with no mention of alcohol consumption patterns, Stockley's described that both NSAIDs and excessive alcohol use carry the risk of gastrointestinal adverse effects, while NSAIDs and alcohol were not included as an interaction in the BNF. For the next step, the Project Steering Group reviewed and assessed each medication or drug class. Using the definition of "necessary to avoid" previously described by Dreischulte et al. (37), it was considered necessary to avoid concurrent alcohol with a specific medication if, in an average older adult concurrent use of alcohol with this medication would be considered unsafe because (a) there is sufficient evidence that the patient is likely to be harmed, and (b) the likely harm to the patient is large enough to be clinically significant.

Drugs were excluded for the following reasons: their interaction with alcohol was not considered serious i.e. unlikely to cause significant harm to the patient (for example: selective serotonin reuptake inhibitors (SSRIs) and alcohol) (33); the medication is not licensed for use in older adults; the medication is only administered in a hospital environment (for example Trabectedin), interaction with alcohol may only occur during alcohol withdrawal or has been withdrawn from the market in the UK or Ireland. Following these exclusions a truncated list was further reviewed by the Project Steering Group by a consensus discussion. In this process the Project Steering Group agreed, that BNF black dot warnings would be included in the final list. BNF black dots refer to potentially serious drug-alcohol interactions where concurrent use should be avoided or only undertaken with caution and appropriate management. In the paper version of the BNF, these potentially serious drug-alcohol interactions are highlighted to prescribers by flagging them with a 'black dot' (31). Furthermore, the amount of alcohol to be avoided with each medication class was defined through group discussion, based on both the evidence available and the members' own clinical experience. Additionally, it was agreed that the following adverse outcomes associated with concurrent use were defined using clinical guidance documents and reviews to ensure clarity among all participants: orthostatic hypotension (38), hypoglycaemia (39) and lactic acidosis (40).

Selection of the Delphi panel

A total of 51 experts from the Republic of Ireland and United Kingdom (UK), were invited (via e-mail or letter) to participate as part of the Delphi consensus panel. The experts invited to participate on the Delphi consensus panel were peer recognized by the Project Steering Group or nominated by other panel members on the basis of their clinical experience or knowledge of alcohol-medication interactions or care of older persons.

Reasons for non-participation were not required; however, in some instances they were provided, they included time commitments and lack of clinical knowledge of alcohol-medication interactions. In total, 19 participants (37%) agreed to participate and written consent was received from all participants before the study. The panel consisted of general practitioners (GPs) (n=5), geriatricians (n=3), hospital pharmacists with expertise in care of older adults (n=3), community pharmacists (n=3), clinical pharmacologists (n=2), clinical pharmacists specialising in substance misuse (n=2) and a physician specialising in substance misuse (n=1). Panel members were not provided with compensation for participation.

The Delphi validation technique and process

An online questionnaire was piloted by two pharmacists and two GPs, to identify any potential problems and to approximate the completion time for the survey. Following amendments, the 19 participants were sent a link to the online POSAMINO questionnaire (via SurveyGizmo®) in March 2016. Participants were given four weeks to complete the online questionnaire and all participants were sent a reminder email after 2 weeks to encourage participation. The questionnaire was both anonymous and confidential.

The quantity of alcohol per standard drink (SD) (10 grammes alcohol in Ireland) or unit (8 grammes alcohol in the United Kingdom), was defined at the beginning of the survey. As some interactions are listed with any alcohol consumption and others with 'heavy' consumption, we provided definitions of both at the beginning of the questionnaire. Any alcohol consumption was defined as ≥ 1 SD of alcohol in Ireland or ≥ 1 unit of alcohol in the UK, with heavy consumption defined according to Irish and UK National low risk drinking guidelines. (35, 36) The panel were instructed to evaluate each potential interaction listed according to the

“necessary to avoid” framework described above. (37) Medications were categorised according to the BNF physiological classification system. Each statement was presented in the same format stating: It is necessary to avoid [quantity of alcohol] (any or heavy) with [medications/ drug class], followed by a brief rationale for the statement. For example: *It is necessary to avoid: Any alcohol consumption with first generation anti-histamines (for example promethazine). (Rationale: Concurrent alcohol consumption with first generation anti-histamines can increase the risk of sedation).*

Participants were asked to rate their agreement with each statement, using a 5-point Likert scale (1-strongly disagree to 5-strongly agree) along with any additional comments, or suggestions for additional medicines to be included.

After each round, the median and interquartile ranges (IQR) were calculated for each statement. Consistent with previous Delphi consensus studies, the consensus level required for a statement to be retained was defined *a priori* as a median of 4 or 5 with a lower quartile value of ≥ 4 . (41) If a statement had an upper quartile value of ≤ 2 , this indicated there was general disagreement with the statement between panel members, and the statement was rejected. If group consensus was not reached, the criteria were reviewed by the Project Steering Group and were removed based on comments or revised and included in the second questionnaire. In the second questionnaire, panellists were provided with links to the most recent evidence relating to each of the alcohol-medication interactions, to help facilitate their decisions due to uncertainty in round 1. As with round one, the median response and IQR were calculated and evaluated by the Project Steering Group using the same thresholds to determine consensus between the panel members. If consensus was not reached following the second round, the statement was rejected. Statistical analysis was performed using STATA version 13 and Microsoft Excel 2010.

Results

Following an initial review of the literature, Anatomical Therapeutic Chemical (ATC) codes for a total of 364 AI medicines were extracted by the Project Steering Group (**Supplementary File 1**). Medicines were classified according to drug classes and organised according to BNF physiological classification systems. From this list, a total of 63 statements were initially compiled based on the 'necessary to avoid' framework. (37) Following group discussions, 11 statements were removed/updated or merged together, based on the steering groups clinical knowledge or experience with medicines. For example, the steering group decided to no longer classify benzodiazepines according to the duration of action, as both long acting and short acting have the potential to interact with alcohol. Furthermore, a total of eight BNF black dot warnings were included in the final criteria as they represent potentially serious interactions.

All 19 panel members completed the round one questionnaire, in which forty-four statements were presented. Consensus was reached on 13 statements, with no statements rejected (**Table 1**). Thirty-one statements were reviewed by the steering group, with nine statements removed based on comments from the panel, if the interaction was not relevant to older adults or the interaction was not of clinical significance for example: heavy alcohol consumption with vitamin A preparations. Furthermore, eight new statements were included in round two based on comments from panellists at the first round.

A total of 18 out of the 19 recruited participants completed the round two questionnaire. Of the 30 statements included in the second questionnaire, consensus was reached for 17 statements. The remaining 13 statements were rejected as no consensus was reached. In total, consensus was reached for 30 potentially serious drug alcohol interactions in older adults, with the inclusion of BNF black dot warnings the final list was 38 statements.

INSERT TABLE 1 HERE

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

This final 38 item POSAMINO criteria (**Table 2**) were organised over the following physiological systems:
Central nervous system (n=15), cardiovascular system (n=9), endocrine system (n=5), musculoskeletal system
(n=3), infections (n=3), malignant disease and immunosuppression (n=2) and respiratory system (n=1).

INSERT TABLE 2 HERE

For peer review only

Discussion

Principle findings in relation to previous studies

While older adults experience a disproportionate burden of alcohol related harm, (5) research suggests that healthcare professionals are less likely to discuss alcohol consumption with older adults. (42, 43) Flagging older adults at the point of prescribing an alcohol-interactive medication may facilitate targeted screening and interventions to help reduce harm. Despite the high propensity for alcohol-medication interactions among community dwelling older adults, our recent systematic review has highlighted that there is still a lack of consensus regarding what constitutes an AI medication. (30) This study reports the development of a set of 38 explicit criteria, POSAMINO, for identifying older adults at risk of potentially serious alcohol-medication interactions.

The final POSAMINO criteria consist of seven different drug classes, with central nervous system (CNS) medicines representing 40% of the criteria. Estimates from our previous study indicate that approximately one in five older Irish adults are exposed to CNS agents, with over half of patients using CNS agents also reporting concurrent alcohol consumption. (12) Nine of the final 38 criteria concern cardiovascular agents, another widely used drug class among aged adults. (12) There is also a high-risk of adverse effects associated with these agents, with anti-platelets, diuretics and anticoagulants identified as the most common drug classes involved in preventable drug-related admissions, in a previous study. (44) Analgesics such as non-steroidal anti-inflammatory drugs (NSAIDs), opioids, paracetamol and gabapentin (used for neuropathic pain) were also included in the final POSAMINO criteria. The prevalence of chronic pain increases with age and older adults may consume alcohol to help cope and manage their pain. (45) This behaviour may also predispose older adults to adverse outcomes associated with concurrent use with analgesics (for example increased sedation, falls and gastrointestinal bleeds).

The development of the POSAMINO criteria is important for several reasons. In the absence of an explicit list of alcohol-interactive medications, multiple drug reference sources have been used in previous studies leading to a lack of consistency in the inclusion of medications across studies, (11, 12, 19, 22, 23) which may result in biased estimates of potential risk among older adults. Furthermore, some of these interactions may be

theoretical with trivial clinical significance. Therefore, this study developed a consensus-based set of explicit criteria for potentially serious alcohol-medication interactions in older adults rather than an exhaustive list of medications with potential to interact with alcohol. Consistent with existing literature, the POSAMINO criteria classify CNS agents as AI medications with the potential to cause serious harm to older adults. (9-13, 15-24, 26, 28) However, some of the previous studies focussed on sedatives/hypnotics only. (13, 26, 28) We identified a number of additional CNS agents, such as anti-Parkinson's drugs (for example: Pramipexole, Apomorphine and Levodopa), which may be overlooked if we focus solely on sedatives or hypnotics.

Furthermore, the POSAMINO criteria also includes a wide range of other drug classes, such as cardiovascular, respiratory system, musculoskeletal, malignant disease, infections or endocrine agents which were not all considered among previous studies. (9-11, 16, 19, 21, 23) It is also important to note that there has been little emphasis on the adverse outcomes or severity of these potential interactions. (22, 30) The development of the POSAMINO criteria may help identify older adults at risk of potentially serious alcohol-medicine interactions in the future.

Clinical implications and future research

This study adds to a growing body of research investigating the concurrent use of alcohol and alcohol interactive medicines in older adults. Undoubtedly, adverse drug events (ADEs) represent a major burden on healthcare, with ADEs detected in 26.3% of patients aged ≥ 65 years admitted to an Irish hospital with acute illnesses. (46) The increasing use of multiple medications, (6) combined with an increased frequency of alcohol consumption (2) and age-related physiological changes, may predispose older adults to these adverse events. The significant burden of alcohol-related harm and mortality among older adults, (5) indicates a pressing need for future interventions to minimise risk. However, prior to informing clinical or public health initiatives, the final criteria developed from this study, will require further validation to prospectively quantify the magnitude of risk posed for each criterion for adverse outcomes in older adults. In particular, the quantity of alcohol specified in each criterion will need to be further evaluated in prospective studies. Once validated, these criteria have the potential to inform medical decision making and enable healthcare professionals to risk stratify older adults at the point of prescribing and prioritise alcohol screening and brief interventions in high-

1
2
3 risk groups. This is essential, as older people have previously reported little knowledge about risks associated
4
5 with the concurrent use of alcohol and medications. (47)
6
7

8 *Strengths and Limitations*

9

10 The criteria were developed in a robust fashion, using a two-step process involving a systematic review and
11 two-round Delphi process. The Delphi technique is flexible and for this study enabled communication from a
12 group of nineteen healthcare professionals from both the United Kingdom and Ireland. Eighteen of the 19
13 participants completed the two rounds, with participants providing numerous comments and suggestions in
14 both rounds. Similar to previous studies, participants remained anonymous and were not provided with
15 feedback following the first round in order to remove the risk of potential bias. (41)
16
17
18
19
20
21
22

23 Inevitably, there were limitations to this Delphi study. With all Delphi studies, participants' judgement may be
24 subjective. In order to reduce this potential bias, a diverse group of healthcare professionals with expertise or
25 interest in the care of older adults were selected. With explicit criteria, there is also a need for regular updates
26 and revision due to the availability of more up to date information after development. (48) The developed
27 POSAMINO criteria do not apply to older adults diagnosed with an alcohol use disorder, as chronic heavy
28 alcohol consumption can substantially increase the activity of the cytochrome P450 metabolising enzyme
29 CYP2E1. (8) Finally, older adults may also experience chronic illnesses which affect the metabolism of both
30 alcohol and medications. As a result, it is important that healthcare professionals also consider these comorbid
31 diseases when assessing the risk for potential adverse outcomes. (49)
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Conclusions

Using a systematic review and a two-round Delphi consensus method, we have developed the first set of explicit potentially serious alcohol-medication interactions (POSAMINO) for use in older adults. Following future validation studies, these criteria may allow for the risk stratification of older adults at the point of prescribing, and prioritise alcohol screening and brief alcohol interventions in high-risk groups.

For peer review only

Declarations:

Authors' contributions: AH, PG, CR, TF & GC (the project steering group) conceived and designed this study. AH & GC conducted initial literature search. AH communicated with Delphi participants. AH analysed the data after each round. All Project Steering Group members were involved in interpretation of the data. AH & GC drafted the manuscript. All co-authors revised the manuscript and have given the approval for publication.

Competing interests: The authors declare that they have no competing interests.

Ethics approval: Ethical approval for this study was approved by the Royal College of Surgeons in Ireland (RCSI) ethics committee (REC 1097).

Acknowledgements: We would like to thank all Delphi participants who contributed to the development of the POSAMINO criteria.

Funding: This study was funded by the Royal College of Surgeons in Ireland (RCSI), School of Pharmacy, Dublin 2, Ireland.

Data sharing statement: Additional data is available by request from the corresponding author.

References

1. United Nations, Department of Economic and Social Affairs PD. World Population Prospects: The 2015 Revision, Key Findings and Advance Tables. New York: 2015. Available from: https://esa.un.org/unpd/wpp/Publications/Files/Key_Findings_WPP_2015.pdf

2. Britton A, Ben-Shlomo Y, Benzeval M, Kuh D, Bell S. Life course trajectories of alcohol consumption in the United Kingdom using longitudinal data from nine cohort studies. *BMC Med.* 2015;13:47.

3. Meng Y, Holmes J, Hill-McManus D, Brennan A, Meier PS. Trend analysis and modelling of gender-specific age, period and birth cohort effects on alcohol abstinence and consumption level for drinkers in Great Britain using the General Lifestyle Survey 1984–2009. *Addiction.* 2014;109.

4. Huang AR, Mallet L, Rochefort CM, Egual T, Buckeridge DL, Tamblyn R. Medication-related falls in the elderly: causative factors and preventive strategies. *Drugs Aging.* 2012;29(5):359-76.

5. Wadd S, Papadopoulos C. Drinking behaviour and alcohol-related harm amongst older adults: analysis of existing UK datasets. *BMC Res Notes.* 2014;7:741.

6. Moriarty F, Hardy C, Bennett K, Smith SM, Fahey T. Trends and interaction of polypharmacy and potentially inappropriate prescribing in primary care over 15 years in Ireland: a repeated cross-sectional study. *BMJ open.* 2015;5.

7. Moore AA, Whiteman EJ, Ward KT. Risks of combined alcohol/medication use in older adults. *Am J Geriatr Pharmacother.* 2007;5.

8. Weathermon R, Crabb DW. Alcohol and medication interactions. *Alcohol research & health : the journal of the National Institute on Alcohol Abuse and Alcoholism.* 1999;23(1):40-54.

9. Adams WL. Potential for adverse drug-alcohol interactions among retirement community residents. *J Am Geriatr Soc.* 1995;43.

10. Aira M, Hartikainen S, Sulkava R. Community prevalence of alcohol use and concomitant use of medication - A source of possible risk in the elderly aged 75 and older? *Int J Geriatr Psychiatry.* 2005;20(7):680-5.

11. Breslow RA, Dong C, White A. Prevalence of alcohol-interactive prescription medication use among current drinkers: United States, 1999 to 2010. *Alcohol Clin Exp Res.* 2015;39(2):371-9.

12. Cousins G, Galvin R, Flood M, Kennedy M-C, Motterlini N, Henman MC, et al. Potential for alcohol and drug interactions in older adults: evidence from the Irish longitudinal study on ageing. *BMC Geriatr.* 2014;14.

13. Del Rio MC, Prada C, Alvarez FJ. Do Spanish patients drink alcohol while undergoing treatment with benzodiazepines? *Alcohol.* 2002;26(1):31-4.

14. Del Río MC, Prada C, Alvarez FJ. The use of medication and alcohol among the Spanish population. *Br J Clin Pharmacol.* 1996;41(3):253-5.

15. Du Y, Scheidt-Nave C, Knopf H. Use of psychotropic drugs and alcohol among non-institutionalised elderly adults in Germany. *Pharmacopsychiatry.* 2008;41(6):242-51.

16. Forster LE, Pollow R, Stoller EP. Alcohol use and potential risk for alcohol-related adverse drug reactions among community-based elderly. *J Community Health*. 1993;18(4):225-39.
17. Ilomaki J, Korhonen MJ, Enlund H, Hartzema AG, Kauhanen J. Risk drinking behavior among psychotropic drug users in an aging Finnish population: the FinDrink study. *Alcohol*. 2008;42(4):261-7.
18. Ilomaeki J, Gnjjidic D, Hilmer SN, Le Couteur DG, Naganathan V, Cumming RG, et al. Psychotropic drug use and alcohol drinking in community-dwelling older Australian men: the CHAMP study. *Drug and alcohol review*. 2013;32(2):218-22.
19. Immonen S, Pitkälä K. The prevalence of potential alcohol-drug interactions in older adults. *Eur Geriatr Med*. 2012;3:S134.
20. John U, Baumeister SE, Volzke H, Meyer C, Ulbricht S, Alte D. Sedative, hypnotic, anxiolytic and opioid medicament use and its co-occurrence with tobacco smoking and alcohol risk drinking in a community sample. *BMC Public Health*. 2007;7:337.
21. Onder G, Landi F, Della Vedova C, Atkinson H, Pedone C, Cesari M, et al. Moderate alcohol consumption and adverse drug reactions among older adults. *Pharmacoepidemiol Drug Saf*. 2002;11(5):385-92.
22. Pringle KE, Ahern FM, Heller DA, Gold CH, Brown TV. Potential for alcohol and prescription drug interactions in older people. *J Am Geriatr Soc*. 2005;53.
23. Qato DM, Manzoor BS, Lee TA. Drug-alcohol interactions in older US adults. *J Am Geriatr Soc*. 2015;63.
24. Sheahan SL, Coons SJ, Robbins CA, Martin SS, Hendricks J, Latimer M. Psychoactive Medication, Alcohol-Use, and Falls among Older Adults. *J Behav Med*. 1995;18(2):127-40.
25. Swift W, Stollznow N, Pirota M. The use of alcohol and medicines among Australian adults. *Aust N Z J Public Health*. 2007;31.
26. Veldhuizen S, Wade TJ, Cairney J. Alcohol consumption among Canadians taking benzodiazepines and related drugs. *Pharmacoepidemiol Drug Saf*. 2009;18.
27. Wong H, Heuberger R, Logomarsino J, Hewlings S. Associations between alcohol use, polypharmacy and falls in older adults. *Nurs Older People*. 2016;28(1):30-6.
28. Lagnaoui R, Moore N, Dartigues JF, Fourrier A, Begaud B. Benzodiazepine use and wine consumption in the French elderly. *Br J Clin Pharmacol*. 2001;52(4):455-6.
29. Hsu C-C, Sandford BA. The Delphi technique: making sense of consensus. *Practical assessment, research & evaluation*. 12(10):1-8.
30. Holton AE, Gallagher P, Fahey T, Cousins G. Concurrent use of alcohol interactive medications and alcohol in older adults: a systematic review of prevalence and associated adverse outcomes. *BMC Geriatr*. 2017;17(1):148.
31. Joint Formulary Committee (2015) The British National Formulary (BNF). 69th ed. London-British Medical Association and Royal Pharmaceutical Society of Great Britain
32. Sweetman S. Martindale: The Complete Drug Reference [online] London: Pharmaceutical Press; Available from: <http://www.medicinescomplete.com/>

33. Stockley's Drug Interactions [online] [Internet]. Pharmaceutical Press. Available from: <https://www.medicinescomplete.com/mc/stockley/current/>.

34. National Institute on Alcohol Abuse and Alcoholism (NIAAA). Harmful Interactions mixing alcohol with medicines. 2014. Available from: https://pubs.niaaa.nih.gov/publications/medicine/Harmful_Interactions.pdf

35. Hope. A Standard Drink in Ireland: What strength? Health Service Executive- Alcohol Implementation Group 2009

36. United Kingdom (UK) Chief Medical Officers' Alcohol Guidelines Review Summary of the proposed new guidelines. Department of Health January 2016 Available from: https://pubs.niaaa.nih.gov/publications/medicine/Harmful_Interactions.pdf

37. Dreischulte T, Grant AM, McCowan C, McAnaw JJ, Guthrie B. Quality and safety of medication use in primary care: consensus validation of a new set of explicit medication assessment criteria and prioritisation of topics for improvement. BMC Clin Pharmacol. 2012;12:5.

38. Lanier JB, Mote MB, Clay EC. Evaluation and Management of Orthostatic Hypotension. Am Fam Physician. 2011;84(5):527-36.

39. European Medicines Agency (EMA). Guideline on clinical investigation of medicinal products in the treatment or prevention of diabetes mellitus Committee for Medicinal Products for Human Use, 2012 Available from: http://www.ema.europa.eu/docs/en_GB/document_library/Scientific_guideline/2012/06/WC500129256.pdf

40. Salpeter SR, Greyber E, Pasternak GA, Salpeter EE. Risk of fatal and nonfatal lactic acidosis with metformin use in type 2 diabetes mellitus: Systematic review and meta-analysis. Arch Intern Med. 2003;163(21):2594-602.

41. Cooper JA, Ryan C, Smith SM, Wallace E, Bennett K, Cahir C, et al. The development of the PROMPT (PRescribing Optimally in Middle-aged People's Treatments) criteria. BMC Health Serv Res. 2014;14:484.

42. Duru OK, Xu H, Tseng C-H, Mirkin M, Ang A, Tallen L, et al. Correlates of Alcohol-Related Discussions between Older Adults and Their Physicians. J Am Geriatr Soc. 2010;58(12):2369-74.

43. Kuerbis A, Sacco P, Blazer DG, Moore AA. Substance Abuse Among Older Adults. Clin Geriatr Med. 2014;30(3):629-54.

44. Howard RL, Avery AJ, Slavenburg S, Royal S, Pipe G, Lucassen P, et al. Which drugs cause preventable admissions to hospital? A systematic review. Br J Clin Pharmacol. 2007;63(2):136-47.

45. Ahangari A, Stewart Williams J, Myleus A. Pain and alcohol consumption among older adults: findings from the World Health Organization Study on global AGEing and adult health, Wave 1. Trop Med Int Health. 2016;21(10):1282-92.

46. Hamilton H, Gallagher P, Ryan C, Byrne S, O'Mahony D. Potentially inappropriate medications defined by stopp criteria and the risk of adverse drug events in older hospitalized patients. Arch Intern Med. 2011;171(11):1013-9.

47. Zanjani F, Hoogland AI, Downer BG. Alcohol and prescription drug safety in older adults. Drug, healthcare and patient safety. 2013;5:13-27.

- 1
2
3
4 48. O'Mahony D, O'Sullivan D, Byrne S, O'Connor MN, Ryan C, Gallagher P. STOPP/START criteria
5 for potentially inappropriate prescribing in older people: version 2. *Age Ageing*. 2015;44(2):213-8.
6
7 49. Slattum PW, Hassan OE. Medications, Alcohol, and Aging. In: Kuerbis A, Moore AA, Sacco P,
8 Zanjani F, editors. *Alcohol and Aging: Clinical and Public Health Perspectives*. Cham: Springer International
9 Publishing; 2016. p. 117-29.
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1: Results from the 2-round Delphi process

	Potentially serious interactions as listed in the BNF	Round 1					Round 2					Final Criteria
Physiological System		Total	Accept	Revise	Reject	Removal	New statements included	Total	Accept	Revise	Reject	
Cardiovascular System	1	14	1	13	0	3	2	12	7	0	5	9
Respiratory System	0	1	1	0	0	0	0	0	0	0	0	1
Central Nervous System	4	12	6	6	0	1	5	10	5	0	5	15
Infections	2	5	0	5	0	3	0	2	1	0	1	3
Endocrine System	0	5	3	2	0	0	0	2	2	0	0	5
Obstetrics, Gynaecology, and Urinary Tract Disorders	0	2	0	2	0	1	1	2	0	0	2	0
Malignant Disease and Immunosuppression	1	1	0	1	0	0	0	1	1	0	0	2
Nutrition and Blood	0	1	0	1	0	1	0	0	0	0	0	0
Musculoskeletal System	0	3	2	1	0	0	0	1	1	0	0	3
Total	8	44	13	31	0	9	8	30	17	0	13	38

Table 2: Final POSAMINO Criteria (n=38)

Cardiovascular System

1. **Heavy alcohol consumption** with multiple anti-hypertensive combinations
Rationale: Concurrent use of alcohol consumption and anti-hypertensives may increase the risk of orthostatic hypotension
2. **Heavy alcohol consumption** with warfarin (& phenindione)
Rationale: Heavy episodic alcohol consumption is associated with an increased risk of major bleeds
3. **Heavy alcohol consumption** with regular use of low dose aspirin (75mg)
Rationale: Heavy alcohol consumption combined with aspirin may cause a small increase in gastrointestinal blood loss
4. **Heavy alcohol consumption** with both regular and as required nitrates (e.g. glyceryl trinitrate, isosorbide dinitrate and isosorbide mononitrate)
Rationale: The combined haemodynamic effects of alcohol and nitrates may increase the risk of exaggerated hypotension.
5. **Heavy alcohol consumption** with the vasodilatory medication nicorandil
Rationale: The combined haemodynamic effects of alcohol & nicorandil may increase the risk of exaggerated hypotension.
6. **Heavy alcohol consumption** with the combined use of both nitrates and vasodilator medication (e.g. nicorandil)
Rationale: The combined haemodynamic effects of alcohol with nitrates and vasodilator drugs, may increase the risk of exaggerated hypotension.
7. **Heavy alcohol consumption** with diuretics (e.g. loop diuretics (furosemide), thiazide diuretics (bendroflumethiazide) & potassium sparing diuretics (amiloride))
Rationale: The concurrent use of alcohol and anti-hypertensives may increase the risk of orthostatic hypotension.
8. **Heavy alcohol consumption** with alpha blockers (e.g. terazosin)
Rationale: The concurrent use of alcohol and anti-hypertensives may increase the risk of orthostatic hypotension.
9. **Heavy alcohol consumption** with centrally acting anti-hypertensives (e.g. clonidine or methyldopa)
Rationale: Alcohol consumption combined with centrally acting anti-hypertensives may increase the risk of sedation and/or orthostatic hypotension

Respiratory System

1. **Any alcohol consumption** with first generation anti-histamines (e.g. promethazine)
Rationale: Concurrent alcohol consumption with first generation anti-histamines can increase the risk of sedation

Central Nervous System (CNS)

1. **Heavy alcohol consumption** with benzodiazepines (e.g. diazepam) and benzodiazepine related medications (e.g. zopiclone)
Rationale: Alcohol consumption combined with benzodiazepines and benzodiazepine related medications may enhance CNS depressant effects
2. **Heavy alcohol consumption** combined with opioids
Rationale: Alcohol consumption combined with opioids may enhance CNS depressant effects of alcohol
3. **Heavy alcohol consumption** with duloxetine
Rationale: Heavy alcohol consumption combined with duloxetine may increase the risk of hepatotoxicity
4. **Heavy alcohol consumption** with all anti-psychotics
Rationale: Alcohol consumption combined with antipsychotics may increase the risk of sedation
5. **Any alcohol consumption** with barbiturates
Rationale: Alcohol consumption combined with barbiturates may increase the risk of sedation.
6. **Heavy alcohol consumption** with anti-epileptic drugs (AEDs)
Rationale: Heavy alcohol consumption can increase the risk of seizures and sedation in patients taking AEDs.
7. **Any alcohol consumption** with tricyclic anti-depressants (TCAs)
Rationale: Alcohol consumption combined with antidepressants may enhance the CNS depressant effects of alcohol
8. **Any alcohol consumption** with tetracyclic antidepressants
Rationale: Alcohol consumption combined with antidepressants may enhance the CNS depressant effects of alcohol
9. **Any alcohol consumption** with mirtazapine
Rationale: Alcohol consumption combined with antidepressants may enhance the CNS depressant effects of alcohol
10. **Any alcohol consumption** with monoamine oxidase inhibitors (MAOIs)
Rationale: A potentially life-threatening hypertensive reaction can develop in patients taking non-selective MAOIs who consume drinks rich in tyramine (e.g. wines, beers and non-alcoholic beers)
11. **Heavy alcohol consumption** with long term regular paracetamol use (e.g. 1g four times a day)
Rationale: Heavy alcohol consumption may increase the risk of paracetamol hepatotoxicity especially if alcohol intake is abruptly stopped.
12. **Heavy alcohol consumption** with gabapentin (when used for neuropathic pain)
Rationale: Alcohol combined with medications for neuropathic pain may increase the risk of sedation
13. **Heavy alcohol consumption** with pramipexole or amantadine
Rationale: Alcohol combined with pramipexole or amantadine may increase the risk of sedation
14. **Heavy alcohol consumption** with apomorphine
Rationale: Alcohol combined with apomorphine may increase the risk of orthostatic hypotension

15. **Heavy alcohol consumption** with levodopa (alone or in combination with carbidopa)
Rationale: Alcohol combined with levodopa (alone or in combination with carbidopa) may increase the risk of orthostatic hypotension

Endocrine

1. **Heavy alcohol consumption** with insulin
Rationale: Alcohol consumption may enhance the hypoglycaemic effect of insulin
2. **Heavy alcohol consumption** with metformin
Rationale: Heavy alcohol consumption combined with metformin may increase the risk of lactic acidosis
3. **Heavy alcohol consumption** with sulphonylureas
Rationale: Alcohol consumption can enhance the hypoglycaemic effects of antidiabetics
4. **Heavy alcohol consumption** with meglitinides (e.g. nateglinide)
Rationale: Alcohol consumption can enhance the hypoglycaemic effects of antidiabetics
5. **Heavy alcohol consumption** with thiazolidinediones (e.g. pioglitazone)
Rationale: Alcohol consumption can enhance the hypoglycaemic effects of antidiabetics

Musculoskeletal and joint diseases

1. **Heavy alcohol consumption** with any nonsteroidal anti-inflammatory drugs (NSAIDs) (including COX-2 inhibitors)
Rationale: Heavy alcohol consumption and NSAID use carry an increased risk of gastrointestinal bleeds
2. **Heavy alcohol consumption** combined with methotrexate or leflunomide
Rationale: Heavy alcohol consumption combined with methotrexate or leflunomide may increase the risk of hepatotoxicity
3. **Heavy alcohol consumption** with oral muscle relaxants (e.g. baclofen)
Rationale: Concurrent alcohol consumption and muscle relaxants can increase the risk of CNS depression

Malignant disease and immunosuppression

1. **Any alcohol consumption** with procarbazine
Rationale: A disulfiram-like reaction can occur when alcohol is given with procarbazine
2. **Heavy alcohol consumption** with interferon alpha or interferon beta
Rationale: Heavy alcohol consumption combined with interferons may increase the risk of hepatotoxicity and reduce the response to treatment with interferon

Infections

1. **Heavy alcohol consumption** with anti-mycobacterial medications such as isoniazid, pyrazinamide, ethionamide and rifampicin (alone or in combination)
Rationale: Alcohol combined with anti-mycobacterial medications can increase the risk of hepatotoxicity
2. **Any alcohol consumption** with cycloserine
Rationale: Alcohol consumption may increase the risk of seizures in patients taking cycloserine
3. **Any alcohol consumption** with metronidazole or tinidazole
Rationale: A disulfiram-like reaction can occur when alcohol is given with metronidazole




Figure 1: Flow chart summarizing the steps involved in the development of the POSAMINO criteria